

CORRECTED VERSION

(19) World Intellectual Property Organization  
International Bureau(43) International Publication Date  
31 January 2002 (31.01.2002)

PCT

(10) International Publication Number  
WO 02/008187 A1(51) International Patent Classification: C07D 209/02,  
211/04, 233/56, 317/10, 319/04, 339/02, 339/08, C07C  
229/00, 233/05, 271/08, 271/32, A61K 31/16, 31/27,  
31/195, 31/357, 31/385, 31/403, 31/445, 31/4164

(21) International Application Number: PCT/US01/22813

(22) International Filing Date: 19 July 2001 (19.07.2001)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
60/220,107 21 July 2000 (21.07.2000) US(71) Applicants: SCHERING CORPORATION [US/US];  
Patent Department K-6-1 1990, 2000 Galloping Hill Road,  
Kenilworth, NJ 07033-0530 (US). CORVAS INTERNA-  
TIONAL, INC. [US/US]; 3030 Science Park Road, San  
Diego, CA 92121 (US).(72) Inventors: SAKSENA, Anil, K.; 53 Beverley Road,  
Upper Montclair, NJ 07059 (US). GIRIJAVALLABHAN,  
Viyyoor, Moopit; 10 Maplewood Drive, Parsippany, NJ  
07054 (US). BOGEN, Stephane, L.; 13 Dahlia Road,  
Somerset, NJ 08873 (US). LOVEY, Raymond, G.; 65  
Woodside Avenue, West Caldwell, NJ 07006 (US). JAO,  
Edwin, E.; 20 Crosswood Way, Warren, NJ 07059 (US).  
BENNETT, Frank; 419 Draco Road, Piscataway, NJ  
08854 (US). MC CORMICK, Jinping, L.; 5 Pace Drive,  
Edison, NJ 08820 (US). WANG, Haiyan; 5 Cubberly  
Court, Cranbury, NJ 08512 (US). PIKE, Russell, E.;  
RD #1, 31 Florence Street, Stanhope, NJ 07874 (US).  
LIU, Yi-Tsung; 34 Alexandria Road, Morris Township,  
NJ 07960 (US). CHAN, Tin-Yau; 26 Barlow Road,  
Edison, NJ 08817 (US). ZHOU, Zhaouing; 34 Stonehedge  
Drive, East Windsor, NJ 08520 (US). ARASAPPAN,  
Ashok; 18 Larsen Court, Bridgewater, NJ 08807 (US).  
CHEN, Kevin, X.; 44 Gill Lane, Apt. 1D, Iselin, NJ  
08830 (US). VENKATRAMAN, Srikanth; 6 Tulip  
Drive, #3H, Fords, NJ 08863 (US). PAREKH, Tejal,  
N.; 1885 Ednamary Way, Unit C, Mountain View, CA  
94040 (US). PINTO, Patrick, A.; 34 Battle Ridge Road,  
Morris Plains, NJ 07950 (US). SANTHANAM, Bama;10 Somerset Avenue, Bridgewater, NJ 08807 (US).  
NJORGE, F., George; 11 Softwood Way, Warren, NJ  
07059 (US). GANGULY, Ashit, K.; 96 Cooper Avenue,  
Upper Montclair, NJ 07043 (US). VACCARO, Henry, A.;  
123 Somerset Avenue, South Plainfield, NJ 07080 (US).  
KEMP, Scott, Jeffrey; 7873 Avenida Navidad #263,  
San Diego, CA 92122 (US). LEVY, Odile, Esther; 5304  
Ruelle de Mer, San Diego, CA 92130 (US). LIM-WILBY,  
Marguerita; 6333 Castejon Drive, La Jolla, CA 92037  
(US). TAMURA, Susan, Y.; 2213 Calle Cacique, Santa  
Fe, NM 87505 (US).(81) Designated States (national): AF, AG, AI, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CZ,  
DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, HR, HU,  
ID, IL, IN, IS, JP, KG, KR, KZ, LC, LK, LR, LT, LU, LV,  
MA, MD, MG, MK, MN, MX, MZ, NO, NZ, PL, PT, RO,  
RU, SE, SG, SI, SK, SL, TJ, TM, TR, TT, TZ, UA, UZ, VN,  
YU, ZA.(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF,  
CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD,  
TG).

## Declaration under Rule 4.17:

— as to the applicant's entitlement to claim the priority of the  
earlier application (Rule 4.17(iii)) for all designations

## Published:

— with international search report

(48) Date of publication of this corrected version:

3 January 2003

## (15) Information about Correction:

see PCT Gazette No. 01/2003 of 3 January 2003, Section II

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: NOVEL PEPTIDES AS NS3-SERINE PROTEASE INHIBITORS OF HEPATITIS C VIRUS

(57) Abstract: The present invention discloses novel peptide compounds which have HCV protease inhibitory activity as well as methods for preparing such compounds. In another embodiment, the invention discloses pharmaceutical compositions comprising such compounds as well as methods of using them to treat disorders associated with the HCV protease.

WO 02/008187 A1

## **Novel Peptides as NS3-Serine Protease Inhibitors of Hepatitis C Virus**

### **Field of Invention**

The present invention relates to novel hepatitis C virus ("HCV") protease inhibitors, pharmaceutical compositions containing one or more such inhibitors, methods of preparing such inhibitors and methods of using such inhibitors to treat hepatitis C and related disorders. This invention specifically discloses novel peptide compounds as inhibitors of the HCV NS3/NS4a serine protease.

### **Background of the Invention**

Hepatitis C virus (HCV) is a (+)-sense single-stranded RNA virus that has been implicated as the major causative agent in non-A, non-B hepatitis (NANBH), particularly in blood-associated NANBH (BB-NANBH)(see, International Patent Application Publication No. WO 89/04669 and European Patent Application Publication No. EP 381 216). NANBH is to be distinguished from other types of viral-induced liver disease, such as hepatitis A virus (HAV), hepatitis B virus (HBV), delta hepatitis virus (HDV), cytomegalovirus (CMV) and Epstein-Barr virus (EBV), as well as from other forms of liver disease such as alcoholism and primary biliar cirrhosis.

Recently, an HCV protease necessary for polypeptide processing and viral replication has been identified, cloned and expressed; (see, e.g., U.S. Patent No. 5,712,145). This approximately 3000 amino acid polyprotein contains, from the amino terminus to the carboxy terminus, a nucleocapsid protein (C), envelope proteins (E1 and E2) and several non-structural proteins (NS1, 2, 3, 4a, 5a and 5b). NS3 is an approximately 68 kda protein, encoded by approximately 1893 nucleotides of the HCV genome, and has two distinct domains: (a) a serine protease domain consisting of approximately 200 of the N-terminal amino acids; and (b) an RNA-dependent ATPase domain at the C-

terminus of the protein. The NS3 protease is considered a member of the chymotrypsin family because of similarities in protein sequence, overall three-dimensional structure and mechanism of catalysis. Other chymotrypsin-like enzymes are elastase, factor Xa, thrombin, trypsin, plasmin, urokinase, tPA and PSA. The HCV NS3 serine protease is responsible for proteolysis of the polypeptide (polyprotein) at the NS3/NS4a, NS4a/NS4b, NS4b/NS5a and NS5a/NS5b junctions and is thus responsible for generating four viral proteins during viral replication. This has made the HCV NS3 serine protease an attractive target for antiviral chemotherapy.

It has been determined that the NS4a protein, an approximately 6 kda polypeptide, is a co-factor for the serine protease activity of NS3. Autocleavage of the NS3/NS4a junction by the NS3/NS4a serine protease occurs intramolecularly (*i.e.*, *cis*) while the other cleavage sites are processed intermolecularly (*i.e.*, *trans*).

Analysis of the natural cleavage sites for HCV protease revealed the presence of cysteine at P1 and serine at P1' and that these residues are strictly conserved in the NS4a/NS4b, NS4b/NS5a and NS5a/NS5b junctions. The NS3/NS4a junction contains a threonine at P1 and a serine at P1'. The Cys→Thr substitution at NS3/NS4a is postulated to account for the requirement of *cis* rather than *trans* processing at this junction. See, *e.g.*, Pizzi *et al.* (1994) *Proc. Natl. Acad. Sci (USA)* 91:888-892, Failla *et al.* (1996) *Folding & Design* 1:35-42. The NS3/NS4a cleavage site is also more tolerant of mutagenesis than the other sites. See, *e.g.*, Kollykhalov *et al.* (1994) *J. Virol.* 68:7525-7533. It has also been found that acidic residues in the region upstream of the cleavage site are required for efficient cleavage. See, *e.g.*, Komoda *et al.* (1994) *J. Virol.* 68:7351-7357.

Inhibitors of HCV protease that have been reported include antioxidants (see, International Patent Application Publication No. WO 98/14181), certain peptides and peptide analogs (see, International Patent Application Publication No. WO 98/17679, Landro *et al.* (1997) *Biochem.*


36:9340-9348, Ingallinella *et al.* (1998) *Biochem.* 37:8906-8914, Llinàs-Brunet *et al.* (1998) *Bioorg. Med. Chem. Lett.* 8:1713-1718), inhibitors based on the 70-amino acid polypeptide eglin c (Martin *et al.* (1998) *Biochem.* 37:11459-11468, inhibitors affinity selected from human pancreatic secretory trypsin inhibitor (hPSTI-C3) and minibody repertoires (MBip) (Dimasi *et al.* (1997) *J. Virol.* 71:7461-7469), cV<sub>H</sub>E2 (a "camelized" variable domain antibody fragment) (Martin *et al.* (1997) *Protein Eng.* 10:607-614), and  $\alpha$ 1-antichymotrypsin (ACT) (Elzouki *et al.* (1997) *J. Hepat.* 27:42-28). A ribozyme designed to selectively destroy hepatitis C virus RNA has recently been disclosed (see, *BioWorld Today* 9(217): 4 (November 10, 1998)).

Reference is also made to the PCT Publications, No. WO 98/17679, published April 30, 1998 (Vertex Pharmaceuticals Incorporated); WO 98/22496, published May 28, 1998 (F. Hoffmann-La Roche AG); and WO 99/07734, published February 18, 1999 (Boehringer Ingelheim Canada Ltd.).

HCV has been implicated in cirrhosis of the liver and in induction of hepatocellular carcinoma. The prognosis for patients suffering from HCV infection is currently poor. HCV infection is more difficult to treat than other forms of hepatitis due to the lack of immunity or remission associated with HCV infection. Current data indicates a less than 50% survival rate at four years post cirrhosis diagnosis. Patients diagnosed with localized resectable hepatocellular carcinoma have a five-year survival rate of 10-30%, whereas those with localized unresectable hepatocellular carcinoma have a five-year survival rate of less than 1%.


Reference is made to A. Marchetti *et al.*, *Synlett*, S1, 1000-1002 (1999) describing the synthesis of bicyclic analogs of an inhibitor of HCV NS3 protease. A compound disclosed therein has the formula:

4




Reference is also made to W. Han *et al*, *Bioorganic & Medicinal Chem. Lett.* (2000) 10, 711-713, which describes the preparation of certain  $\alpha$ -ketoamides,  $\alpha$ -ketoesters and  $\alpha$ -diketones containing allyl and ethyl functionalities.

Reference is also made to WO 00/09558 (Assignee: Boehringer Ingelheim Limited; Published February 24, 2000) which discloses peptide derivatives of the formula:




where the various elements are defined therein. An illustrative compound of that series is:

5




Reference is also made to WO 00/09543 (Assignee: Boehringer  
 Ingelheim Limited; Published February 24, 2000) which discloses peptide  
 5 derivatives of the formula:



where the various elements are defined therein. An illustrative compound of  
 that series is:

6



Current therapies for hepatitis C include interferon- $\alpha$  (INF $\alpha$ ) and combination therapy with ribavirin and interferon. See, e.g., Beremguer et al. (1998) Proc. Assoc. Am. Physicians 110(2):98-112. These therapies suffer from a low sustained response rate and frequent side effects. See, e.g., Hoofnagle et al. (1997) N. Engl. J. Med. 336:347. Currently, no vaccine is available for HCV infection.

Pending and copending U. S. patent applications, Serial No. 09/-----, filed -----, and Serial No. 09/-----, filed -----, Serial No. 09/-----, filed -----, Serial No. 09/-----, filed -----, Serial No. 09/-----, filed -----, and Serial No. 09/-----, filed -----, disclose various types of peptides as NS-3 serine protease inhibitors of hepatitis C virus.

There is a need for new treatments and therapies for HCV infection. It is, therefore, an object of this invention to provide compounds useful in the treatment or prevention or amelioration of one or more symptoms of hepatitis C.

It is a further object herein to provide methods of treatment or prevention or amelioration of one or more symptoms of hepatitis C.

A still further object of the present invention is to provide methods for modulating the activity of serine proteases, particularly the HCV NS3/NS4a serine protease, using the compounds provided herein.


Another object herein is to provide methods of modulating the  
5 processing of the HCV polypeptide using the compounds provided herein.

### Summary of the invention

In its many embodiments, the present invention provides a novel class of inhibitors of the HCV protease, pharmaceutical compositions containing  
10 one or more of the compounds, methods of preparing pharmaceutical formulations comprising one or more such compounds, and methods of treatment, prevention or amelioration or one or more of the symptoms of hepatitis C. Also provided are methods of modulating the interaction of an HCV polypeptide with HCV protease. Among the compounds provided  
15 herein, compounds that inhibit HCV NS3/NS4a serine protease activity are preferred. The present application discloses peptide compounds containing an amino acid arrangement from P3 up to P2'.

In its first embodiment, the present invention provides a compound of  
Formula I:

20



Formula I

wherein:

25 G, J and Y are independently selected from the moieties: H, alkyl, alkyl-aryl, heteroalkyl, heteroaryl, aryl-heteroaryl, alkyl-heteroaryl, cycloalkyl, alkyloxy,



alkyl-aryloxy, aryloxy, heteroaryloxy, heterocycloalkyloxy, cycloalkyloxy, alkylamino, arylamino, alkyl-aryl amino, arylamino, heteroaryl amino, cycloalkyl amino or heterocycloalkyl amino with the proviso that Y maybe optionally substituted with  $X^{11}$  or  $X^{12}$ ;

- 5  $X^{11}$  is alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkyl-alkyl, heterocycl, heterocyclalkyl, aryl, alkylaryl, arylalkyl, heteroaryl, alkylheteroaryl, or heteroarylalkyl, with the proviso that  $X^{11}$  may be additionally optionally substituted with  $X^{12}$ ;

- $X^{12}$  is hydroxy, alkoxy, aryloxy, thio, alkylthio, arylthio, amino, alkylamino, 10 arylamino, alkylsulfonyl, arylsulfonyl, alkylsulfonamido, arylsulfonamido, carboxy, carbalkoxy, carboxamido, alkoxycarbonylamino, alkoxycarbonyloxy, alkylureido, arylureido, halogen, cyano, or nitro, with the proviso that the alkyl, alkoxy, and aryl may be additionally optionally substituted with moieties selected from  $X^{12}$ ;

- 15  $R^1$  is  $COR^5$  or  $B(OR)^2$ , wherein  $R^5$  is H, OH,  $OR^8$ ,  $NR^9R^{10}$ ,  $CF_3$ ,  $C_2F_5$ ,  $C_3F_7$ ,  $CF_2R^6$ ,  $R^6$ , or  $COR^7$  wherein  $R^7$  is H, OH,  $OR^8$ ,  $CHR^9R^{10}$ , or  $NR^9R^{10}$ , wherein  $R^6$ ,  $R^8$ ,  $R^9$  and  $R^{10}$  are independently selected from the group consisting of H, alkyl, aryl, heteroalkyl, heteroaryl, cycloalkyl, cycloalkyl, arylalkyl, heteroarylalkyl,  $CH(R^{1'})COOR^{11}$ ,  
 20  $CH(R^{1'})CONR^{12}R^{13}$ ,  
 $CH(R^{1'})CONHCH(R^{2'})COOR^{11}$ ,  
 $CH(R^{1'})CONHCH(R^{2'})CONR^{12}R^{13}$ ,  $CH(R^{1'})CONHCH(R^{2'})R^1$ ,  
 $CH(R^{1'})CONHCH(R^{2'})CONHCH(R^{3'})COOR^{11}$ ,  
 $CH(R^{1'})CONHCH(R^{2'})CONHCH(R^{3'})CONR^{12}R^{13}$ ,  
 25  $CH(R^{1'})CONHCH(R^{2'})CONHCH(R^{3'})CONHCH(R^{4'})COOR^{11}$ ,

CH(R<sup>1'</sup>)CONHCH(R<sup>2'</sup>)CONHCH(R<sup>3'</sup>)CONHCH(R<sup>4'</sup>)CONR<sup>12</sup>R<sup>13</sup>,  
 CH(R<sup>1'</sup>)CONHCH(R<sup>2'</sup>)CONHCH(R<sup>3'</sup>)CONHCH(R<sup>4'</sup>)CONHCH(R<sup>5'</sup>)COO-  
 R<sup>11</sup>, and

CH(R<sup>1'</sup>)CONHCH(R<sup>2'</sup>)CONHCH(R<sup>3'</sup>)CONHCH(R<sup>4'</sup>)CONHCH(R<sup>5'</sup>)  
 5 CONR<sup>12</sup>R<sup>13</sup>, wherein R<sup>1'</sup>, R<sup>2'</sup>, R<sup>3'</sup>, R<sup>4'</sup>, R<sup>5'</sup>, R<sup>11</sup>, R<sup>12</sup>, R<sup>13</sup>, and R' are independently selected from the group consisting of H, alkyl, aryl, heteroalkyl, heteroaryl, cycloalkyl, alkyl-aryl, alkyl-heteroaryl, aryl-alkyl and heteroaralkyl;

Z is selected from O, N, or CH;

10 W maybe present or absent, and if W is present, W is selected from C=O, C=S, or SO<sub>2</sub>; and

R, R<sup>2</sup>, R<sup>3</sup> and R<sup>4</sup> are independently selected from the group consisting of H; C1-C10 alkyl; C2-C10 alkenyl; C3-C8 cycloalkyl; C3-C8 heterocycloalkyl, alkoxy, aryloxy, alkylthio, arylthio, amino, amido, ester, carboxylic acid,  
 15 carbamate, urea, ketone, aldehyde, cyano, nitro; oxygen, nitrogen, sulfur, and phosphorus atoms (with said oxygen, nitrogen, sulfur, or phosphorus atoms numbering zero to six);


(cycloalkyl)alkyl and (heterocycloalkyl)alkyl, wherein said cycloalkyl is made of three to eight carbon atoms, and zero to six oxygen, nitrogen, sulfur, or  
 20 phosphorus atoms, and said alkyl is of one to six carbon atoms; aryl; heteroaryl; alkyl-aryl; and alkyl-heteroaryl;

wherein said alkyl, heteroalkyl, alkenyl, heteroalkenyl, aryl, heteroaryl, cycloalkyl and heterocycloalkyl moieties may be optionally substituted, with said term "substituted" referring to optional and chemically-suitable  
 25 substitution with one or more moieties selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, aralkyl, cycloalkyl, heterocyclic, halogen, hydroxy, thio, alkoxy, aryloxy, alkylthio, arylthio, amino, amido, ester, carboxylic acid,

carbamate, urea, ketone, aldehyde, cyano, nitro, sulfonamide, sulfoxide, sulfone, sulfonylurea, hydrazide, and hydroxamate.

Among the above-stated definitions for the various moieties of Formula I, the preferred groups for the various moieties are as follows:

- 5 Preferred definition for  $R^1$  is  $COR^5$  with  $R^5$  being H, OH,  $COOR^8$  or  $CONR^9R^{10}$ , where  $R^8$ ,  $R^9$  and  $R^{10}$  are defined above. Preferred moiety for  $R^1$  is  $COCONR^9R^{10}$ , where  $R^9$  is H and  $R^{10}$  is H,  $CH(R^{1'})COOR^{11}$ ,  $CH(R^{1'})CONR^{12}R^{13}$ ,  $CH(R^{1'})CONHCH(R^{2'})COOR^{11}$ ,  $CH(R^{1'})CONHCH(R^{2'})CONR^{12}R^{13}$  or  $CH(R^{1'})CONHCH(R^{2'})(R')$ . Among these, preferred moieties
- 10 for  $R^{10}$  are:  $CH(R^{1'})CONHCH(R^{2'})COOR^{11}$ ,  $CH(R^{1'})CONHCH(R^{2'})CONR^{12}R^{13}$ ,  $CH(R^{1'})CONHCH(R^{2'})(R')$ , wherein  $R^{1'}$  is H or alkyl, heteroalkyl and  $R^{2'}$  is phenyl, substituted phenyl, hetero atom-substituted phenyl, thiophenyl, cycloalkyl, hetero atom-substituted cycloalkyl, piperidyl and pyridyl. More preferred moieties are: for  $R^{1'}$  is H, and  $R^{11}$  is H or *tert*-butyl;  $R'$  is hydroxymethyl; and
- 15  $R^{2'}$  is selected from the group consisting of:



- 20 wherein:


$U^1$  and  $U^2$  maybe same or different and are selected from the group consisting of H, F,  $CH_2COOH$ ,  $CH_2COOMe$ ,  $CH_2CONH_2$ ,  $CH_2CONHMe$ ,  $CH_2CONMe_2$ , azido, amino, hydroxyl, substituted amino, substituted hydroxyl;

5  $U^3$  and  $U^4$  maybe same or different and are O or S;

$U^5$  is selected from the moieties consisting of alkylsulfonyl, aryl sulfonyl, heteroalkyl sulfonyl, heteroaryl sulfonyl, alkyl carbonyl, aryl carbonyl, heteroalkyl carbonyl, heteroaryl carbonyl, alkoxycarbonyl, aryloxy carbonyl, heteroaryloxy carbonyl, alkylaminocarbonyl, arylaminocarbonyl, heteroarylaminocarbonyl or a combination thereof; and


10  $NR^{12}R^{13}$  is selected from the group consisting of:

$NH_2$ ,  $NHMe$ ,  $N(Me)OMe$ ,  $NMe_2$ ,




wherein  $U^6$  is H, OH, or  $CH_2OH$ .

15 Preferred moieties for  $R^2$  are:



Preferred moieties for  $R^3$  are:




wherein R<sup>31</sup> = OH, or O-alkyl.

5 Additionally,  $R^3$  can also be represented by:


14




wherein  $Y^{19}$  is selected from the following moieties:



5 Still, additionally,  $R^3$  can be represented by:




wherein  $Y^{20}$  is selected from the following moieties:




15

Most preferred moieties for  $R^3$  are:



Additionally, the moiety  $Z-C-R^3$  in Formula I, with  $R^4$  being absent, may be represented by the following structures:




5

Some other preferred moieties are: for Z it is N, for  $R^4$  it is H, and for W it is C=O, or SO<sub>2</sub>. Preferred moieties for Y are:



16




wherein:

$Y^{11}$  is selected from H, COOH, COOEt, OMe, Ph, OPh, NHMe, NHAc, NHPh, CH(Me)<sub>2</sub>, 1-triazolyl, 1-imidazolyl, and NHCH<sub>2</sub>COOH;

$Y^{12}$  is selected from H, COOH, COOMe, OMe, F, Cl, or Br.


Y may also be represented by:



5


wherein:

$Y^{13}$  is selected from the following moieties:



and Y<sup>14</sup> is selected from MeSO<sub>2</sub>, Ac, Boc, *i*Boc, Cbz, or Alloc.


Additional preferred structures for Y are:



wherein

$Y^{15}$  and  $Y^{16}$  may be the same or different and are independently selected from alkyl, aryl or herereoalkyl, or heteroaryl.






wherein:


Y<sup>17</sup> is CF<sub>3</sub>, NO<sub>2</sub>, CONH<sub>2</sub>, OH, COOCH<sub>3</sub>, OCH<sub>3</sub>, OC<sub>6</sub>H<sub>5</sub>, C<sub>6</sub>H<sub>5</sub>, COC<sub>6</sub>H<sub>5</sub>, NH<sub>2</sub>, or COOH;


5 Y<sup>18</sup> is COOCH<sub>3</sub>, NO<sub>2</sub>, N(CH<sub>3</sub>)<sub>2</sub>, F, OCH<sub>3</sub>, CH<sub>2</sub>COOH, COOH, SO<sub>2</sub>NH<sub>2</sub>, or NHCOCH<sub>3</sub>.

Preferred moieties for J are:















Preferred moieties for G are:





Q=



5 Unless defined otherwise, all technical and scientific terms used herein have the same meaning as is commonly understood by one of skill in the art to which this invention belongs. Thus, for example, the term alkyl (including the alkyl portions of alkoxy) refers to a monovalent group derived from a straight or branched chain saturated hydrocarbon by the removal of a single

10 atom having from 1 to 8 carbon atoms, preferably from 1 to 6;

aryl — represents a carbocyclic group having from 6 to 14 carbon atoms and having at least one benzenoid ring, with all available substitutable aromatic carbon atoms of the carbocyclic group being intended as possible



points of attachment. Preferred aryl groups include phenyl, 1-naphthyl, 2-naphthyl and indanyl, and especially phenyl and substituted phenyl;

aralkyl – represents a moiety containing an aryl group linked via a lower alkyl;

5        alkylaryl – represents a moiety containing a lower alkyl linked via an aryl group;

cycloalkyl – represents a saturated carbocyclic ring having from 3 to 8 carbon atoms, preferably 5 or 6, optionally substituted.

heterocyclic – represents, in addition to the heteroaryl groups defined  
10 below, saturated and unsaturated cyclic organic groups having at least one O, S and/or N atom interrupting a carbocyclic ring structure that consists of one ring or two fused rings, wherein each ring is 5-, 6- or 7-membered and may or may not have double bonds that lack delocalized pi electrons, which ring structure has from 2 to 8, preferably from 3 to 6 carbon atoms, e.g., 2- or  
15 3-piperidinyl, 2- or 3-piperazinyl, 2- or 3-morpholinyl, or 2- or 3-thiomorpholinyl;

halogen – represents fluorine, chlorine, bromine and iodine;

heteroaryl – represents a cyclic organic group having at least one O, S and/or N atom interrupting a carbocyclic ring structure and having a sufficient  
20 number of delocalized pi electrons to provide aromatic character, with the aromatic heterocyclyl group having from 2 to 14, preferably 4 or 5 carbon atoms, e.g., 2-, 3- or 4-pyridyl, 2- or 3-furyl, 2- or 3-thienyl, 2-, 4- or 5-thiazolyl, 2- or 4-imidazolyl, 2-, 4- or 5-pyrimidinyl, 2-pyrazinyl, or 3- or 4-pyridazinyl, etc. Preferred heteroaryl groups are 2-, 3- and 4-pyridyl; such heteroaryl  
25 groups may also be optionally substituted. Additionally, unless otherwise specifically defined, the term "substituted or unsubstituted" or "optionally substituted" refers to the subject moiety being optionally and chemically suitably substituted with a moiety belonging to R<sup>12</sup> or R<sup>13</sup>.

Also included in the invention are tautomers, rotamers, enantiomers and other optical isomers of compounds of Formula I, as well as pharmaceutically acceptable salts, solvates and derivatives thereof.

A further feature of the invention is pharmaceutical compositions  
5 containing as active ingredient a compound of Formula I (or its salt, solvate or isomers) together with a pharmaceutically acceptable carrier or excipient.

The invention also provides methods for preparing compounds of Formula I, as well as methods for treating diseases such as, for example, HCV and related disorders. The methods for treating comprise administering  
10 to a patient suffering from said disease or diseases a therapeutically effective amount of a compound of Formula I, or pharmaceutical compositions comprising a compound of Formula I.

Also disclosed is the use of a compound of Formula I for the manufacture of a medicament for treating HCV and related disorders.

15

#### **Detailed description of preferred embodiments**

In one embodiment, the present invention discloses compounds of Formula I as inhibitors of HCV protease, especially the HCV NS3/NS4a serine protease, or a pharmaceutically acceptable derivative thereof, where the  
20 various definitions are given above.

Representative compounds of the invention which exhibit excellent HCV protease inhibitory activity are listed below in **Table 1** along with their activity (ranges of  $K_i^*$  values in nanomolar, nM).

**Table 1: Compounds and HCV protease continuous assay results**

Compound from Example No.	Ki* Range
1	C
2	B
3	C
4	C
5	B
6	C
7	C
8	C
9	C
10	C
11	A
12	C
13	B
14	C
15	B
16	A
17	B
18	A
19	not available
20	C
21	B
22	C
23	C
24	B
25	A
26	C
27	C
28	B
29	C
30	B
31	C
32	B
33	C
34	B
35	C
36	C

37	C
38	B
39	C
40	C
41	B
42	C
43	A
44	C
45	C
47	C
48	B
49	A
50	C
51	B
52	B
53	B
54	B
55	C
56	B
57	B
58	C
59	B
60	C
61	C
62	A
63	C
64	B
65	C
66	B
67	C
68	B
69	B
70	B
71	B
72	B
73	A
74	B
75	B
76	B
77	C
78	B
79	C
80	B
81	B


82	C
83	C
84	C
85	C
86	C
87	B
88	B
89	C
90	B
91	B
92	C
93	C
94	B
95	B
96	A
97	A
98	A
99	B
100	B
101	B
102	B
103	C
104	C
105	B
106	A
107	C
108	B
109	B
110	A
111	B
112	B
113	A
114	A
115	C
116	B
117	B
118	B
119	B
120	A
121	B
122	A
123	C
124	B
125	A

126	A
127	A
128	B
129	C
130	A
131	C
132	B
133	B





HCV continuous assay Ki\* range:

Category A= 0-100; B= 101-1000; C= >1000 nM.


Some of the types of the inventive compounds and methods of  
 5 synthesizing the various types of the inventive compounds of Formula I are  
 listed below, then schematically described, followed by the illustrative  
 Examples.



30



5




10


R = Me

R = Benzyl



31



$X = O^tBu$   
 $X = OH$   
 $X = NH_2$   
 $X = NMeOMe$   
 $X = NMe_2$




$X = O^tBu$   
 $X = OH$



$X = H, Y = tBu; X = tBu, Y = H$




32



R = Propargyl

R = Allyl



5



X= O <sup>t</sup>-butyl

$$X = \text{OH}$$

10




X = O<sup>t</sup>Butyl


$$X = \text{OH}$$

15

33


X = O<sup>t</sup>Butyl

X = OH

X = NMe<sub>2</sub>X = O<sup>t</sup>Bu


X = OH

5

R = <sup>t</sup>Bu

R = H

R = Me






X = H, Y = COOH



X = COOH, Y = H

10

34




5



10

35



5

Depending upon their structure, the compounds of the invention may form pharmaceutically acceptable salts with organic or inorganic acids, or organic or inorganic bases. Examples of suitable acids for such salt formation are hydrochloric, sulfuric, phosphoric, acetic, citric, malonic, salicylic, malic, fumaric, succinic, ascorbic, maleic, methanesulfonic and other mineral and carboxylic acids well known to those skilled in the art. For formation of salts with bases, suitable bases are, for example, NaOH, KOH, NH<sub>4</sub>OH, tetraalkylammonium hydroxide, and the like.

15 In another embodiment, this invention provides pharmaceutical compositions comprising the inventive peptides as an active ingredient. The pharmaceutical compositions generally additionally comprise a

pharmaceutically acceptable carrier diluent, excipient or carrier (collectively referred to herein as carrier materials). Because of their HCV inhibitory activity, such pharmaceutical compositions possess utility in treating hepatitis C and related disorders.

5 In yet another embodiment, the present invention discloses methods for preparing pharmaceutical compositions comprising the inventive compounds as an active ingredient. In the pharmaceutical compositions and methods of the present invention, the active ingredients will typically be administered in admixture with suitable carrier materials suitably selected with  
10 respect to the intended form of administration, i.e. oral tablets, capsules (either solid-filled, semi-solid filled or liquid filled), powders for constitution, oral gels, elixirs, dispersible granules, syrups, suspensions, and the like, and consistent with conventional pharmaceutical practices. For example, for oral administration in the form of tablets or capsules, the active drug component  
15 may be combined with any oral non-toxic pharmaceutically acceptable inert carrier, such as lactose, starch, sucrose, cellulose, magnesium stearate, dicalcium phosphate, calcium sulfate, talc, mannitol, ethyl alcohol (liquid forms) and the like. Moreover, when desired or needed, suitable binders, lubricants, disintegrating agents and coloring agents may also be incorporated  
20 in the mixture. Powders and tablets may be comprised of from about 5 to about 95 percent inventive composition. Suitable binders include starch, gelatin, natural sugars, corn sweeteners, natural and synthetic gums such as acacia, sodium alginate, carboxymethylcellulose, polyethylene glycol and waxes. Among the lubricants there may be mentioned for use in these dosage  
25 forms, boric acid, sodium benzoate, sodium acetate, sodium chloride, and the like. Disintegrants include starch, methylcellulose, guar gum and the like.

Sweetening and flavoring agents and preservatives may also be included where appropriate. Some of the terms noted above, namely disintegrants, diluents, lubricants, binders and the like, are discussed in more  
30 detail below.

Additionally, the compositions of the present invention may be formulated in sustained release form to provide the rate controlled release of any one or more of the components or active ingredients to optimize the therapeutic effects, i.e. HCV inhibitory activity and the like. Suitable dosage forms for sustained release include layered tablets containing layers of varying disintegration rates or controlled release polymeric matrices impregnated with the active components and shaped in tablet form or capsules containing such impregnated or encapsulated porous polymeric matrices.

Liquid form preparations include solutions, suspensions and emulsions. As an example may be mentioned water or water-propylene glycol solutions for parenteral injections or addition of sweeteners and pacifiers for oral solutions, suspensions and emulsions. Liquid form preparations may also include solutions for intranasal administration.

Aerosol preparations suitable for inhalation may include solutions and solids in powder form, which may be in combination with a pharmaceutically acceptable carrier such as inert compressed gas, e.g. nitrogen.

For preparing suppositories, a low melting wax such as a mixture of fatty acid glycerides such as cocoa butter is first melted, and the active ingredient is dispersed homogeneously therein by stirring or similar mixing. The molten homogeneous mixture is then poured into convenient sized molds, allowed to cool and thereby solidify.

Also included are solid form preparations which are intended to be converted, shortly before use, to liquid form preparations for either oral or parenteral administration. Such liquid forms include solutions, suspensions and emulsions.

The compounds of the invention may also be deliverable transdermally. The transdermal compositions may take the form of creams, lotions, aerosols and/or emulsions and can be included in a transdermal patch of the matrix or reservoir type as are conventional in the art for this purpose.

Preferably the compound is administered orally, intravenously or subcutaneously.

Preferably, the pharmaceutical preparation is in a unit dosage form. In such form, the preparation is subdivided into suitably sized unit doses  
5 containing appropriate quantities of the active components, e.g., an effective amount to achieve the desired purpose.

The quantity of the inventive active composition in a unit dose of preparation may be generally varied or adjusted from about 1.0 milligram to about 1,000 milligrams, preferably from about 1.0 to about 950 milligrams,  
10 more preferably from about 1.0 to about 500 milligrams, and typically from about 1 to about 250 milligrams, according to the particular application. The actual dosage employed may be varied depending upon the patient's age, sex, weight and severity of the condition being treated. Such techniques are well known to those skilled in the art.

15 Generally, the human oral dosage form containing the active ingredients can be administered 1 or 2 times per day. The amount and frequency of the administration will be regulated according to the judgment of the attending clinician. A generally recommended daily dosage regimen for oral administration may range from about 1.0 milligram to about 1,000  
20 milligrams per day, in single or divided doses.

Some useful terms are described below:

Capsule - refers to a special container or enclosure made of methyl cellulose, polyvinyl alcohols, or denatured gelatins or starch for holding or containing compositions comprising the active ingredients. Hard shell  
25 capsules are typically made of blends of relatively high gel strength bone and pork skin gelatins. The capsule itself may contain small amounts of dyes, opaquing agents, plasticizers and preservatives.

Tablet- refers to a compressed or molded solid dosage form containing the active ingredients with suitable diluents. The tablet can be prepared by

compression of mixtures or granulations obtained by wet granulation, dry granulation or by compaction.

Oral gel- refers to the active ingredients dispersed or solubilized in a hydrophilic semi-solid matrix.

5 Powder for constitution refers to powder blends containing the active ingredients and suitable diluents which can be suspended in water or juices.

Diluent - refers to substances that usually make up the major portion of the composition or dosage form. Suitable diluents include sugars such as lactose, sucrose, mannitol and sorbitol; starches derived from wheat, corn,  
10 rice and potato; and celluloses such as microcrystalline cellulose. The amount of diluent in the composition can range from about 10 to about 90% by weight of the total composition, preferably from about 25 to about 75%, more preferably from about 30 to about 60% by weight, even more preferably from about 12 to about 60%.

15 Disintegrant - refers to materials added to the composition to help it break apart (disintegrate) and release the medicaments. Suitable disintegrants include starches; "cold water soluble" modified starches such as sodium carboxymethyl starch; natural and synthetic gums such as locust bean, karaya, guar, tragacanth and agar; cellulose derivatives such as  
20 methylcellulose and sodium carboxymethylcellulose; microcrystalline celluloses and cross-linked microcrystalline celluloses such as sodium croscarmellose; alginates such as alginic acid and sodium alginate; clays such as bentonites; and effervescent mixtures. The amount of disintegrant in the composition can range from about 2 to about 15% by weight of the  
25 composition, more preferably from about 4 to about 10% by weight.

Binder - refers to substances that bind or "glue" powders together and make them cohesive by forming granules, thus serving as the "adhesive" in the formulation. Binders add cohesive strength already available in the diluent or bulking agent. Suitable binders include sugars such as sucrose; starches  
30 derived from wheat, corn rice and potato; natural gums such as acacia,



gelatin and tragacanth; derivatives of seaweed such as alginic acid, sodium alginate and ammonium calcium alginate; cellulosic materials such as methylcellulose and sodium carboxymethylcellulose and hydroxypropylmethylcellulose; polyvinylpyrrolidone; and inorganics such as magnesium aluminum silicate. The amount of binder in the composition can range from about 2 to about 20% by weight of the composition, more preferably from about 3 to about 10% by weight, even more preferably from about 3 to about 6% by weight.

Lubricant - refers to a substance added to the dosage form to enable the tablet, granules, etc. after it has been compressed, to release from the mold or die by reducing friction or wear. Suitable lubricants include metallic stearates such as magnesium stearate, calcium stearate or potassium stearate; stearic acid; high melting point waxes; and water soluble lubricants such as sodium chloride, sodium benzoate, sodium acetate, sodium oleate, polyethylene glycols and d,l-leucine. Lubricants are usually added at the very last step before compression, since they must be present on the surfaces of the granules and in between them and the parts of the tablet press. The amount of lubricant in the composition can range from about 0.2 to about 5% by weight of the composition, preferably from about 0.5 to about 2%, more preferably from about 0.3 to about 1.5% by weight.

Glident - material that prevents caking and improve the flow characteristics of granulations, so that flow is smooth and uniform. Suitable glidents include silicon dioxide and talc. The amount of glident in the composition can range from about 0.1% to about 5% by weight of the total composition, preferably from about 0.5 to about 2% by weight.

Coloring agents - excipients that provide coloration to the composition or the dosage form. Such excipients can include food grade dyes and food grade dyes adsorbed onto a suitable adsorbent such as clay or aluminum oxide. The amount of the coloring agent can vary from about 0.1 to about 5% by weight of the composition, preferably from about 0.1 to about 1%.

Bioavailability - refers to the rate and extent to which the active drug ingredient or therapeutic moiety is absorbed into the systemic circulation from an administered dosage form as compared to a standard or control.

Conventional methods for preparing tablets are known. Such methods  
5 include dry methods such as direct compression and compression of granulation produced by compaction, or wet methods or other special procedures. Conventional methods for making other forms for administration such as, for example, capsules, suppositories and the like are also well known.

10 Another embodiment of the invention discloses the use of the pharmaceutical compositions disclosed above for treatment of diseases such as, for example, hepatitis C and the like. The method comprises administering a therapeutically effective amount of the inventive pharmaceutical composition to a patient having such a disease or diseases and in need of such a  
15 treatment.

In yet another embodiment, the compounds of the invention may be used for the treatment of HCV in humans in monotherapy mode or in a combination therapy mode such as, for example, in combination with antiviral agents such as, for example, ribavirin and/or interferon such as, for example,  
20  $\alpha$ -interferon and the like.

As stated earlier, the invention includes tautomers, rotamers, enantiomers and other stereoisomers of the compounds also. Thus, as one skilled in the art appreciates, some of the inventive compounds may exist in suitable isomeric forms. Such variations are contemplated to be within the  
25 scope of the invention.

Another embodiment of the invention discloses a method of making the compounds disclosed herein. The compounds may be prepared by several techniques known in the art. Representative illustrative procedures are outlined in the following reaction schemes. It is to be understood that while the  
30 following illustrative schemes describe the preparation of a few representative

inventive compounds, suitable substitution of any of both the natural and unnatural amino acids will result in the formation of the desired compounds based on such substitution. Such variations are contemplated to be within the scope of the invention.

- 5        Abbreviations which are used in the descriptions of the schemes, preparations and the examples that follow are:
- THF: Tetrahydrofuran  
DMF: *N,N*-Dimethylformamide  
EtOAc: Ethyl acetate  
10    AcOH: Acetic acid  
HOObt: 3-Hydroxy-1,2,3-benzotriazin-4(3*H*)-one  
EDCl: 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride  
DEC: 1-(3-dimethylaminopropyl)-3-ethylcarbodiimide hydrochloride  
NMM: *N*-Methylmorpholine  
15    ADDP: 1,1'-(Azodicarbonyl)dipiperidine  
DEAD: Diethylazodicarboxylate  
MeOH: Methanol  
EtOH: Ethanol  
Et<sub>2</sub>O: Diethyl ether  
20    DMSO: Dimethylsulfoxide  
HOBT: *N*-Hydroxybenzotriazole  
PyBrOP: Bromo-*tris*-pyrrolidinophosphonium hexafluorophosphate  
Bn: Benzyl  
Bzl: Benzyl  
25    Et: Ethyl  
Ph: Phenyl  
iBoc: isobutoxycarbonyl  
iPr: isopropyl  
<sup>t</sup>Bu or Bu<sup>t</sup>: *tert*-Butyl  
30    Boc: *tert*-Butoxycarbonyl

Cbz: Benzyloxycarbonyl

Cp: Cyclopentylidienyl

Ts: p-toluenesulfonyl

Me: Methyl

5 THP: Tetrahydropyranyl

iBoc: isobutyloxycarbonyl


Chg: cyclohexylglycine

**General Preparative Schemes:**


The following schemes describe the methods of synthesis of  
10 intermediate building blocks:

44


SCHEME 1




SCHEME 2



SCHEME 3




SCHEME 4




5

**Preparation of Intermediates:****Preparative Example 1**

Step A:Compound (1.1)

To a stirred solution of compound (1.08) (3.00 g, 12.0 mmol; Harbeson, S. L.; Abelleira, S. M.; Akiyama, A.; Barrett, R.; Carroll, R. M.; et al.; J. Med. Chem.; **37** (18) 1994; 2918-2929;) in DMF (15 mL) and CH<sub>2</sub>Cl<sub>2</sub> (15 mL) at -20°C was added HOObt (1.97 g, 12.0 mmol), *N*-methyl morpholine (4.0 mL, 36.0 mmol) and EDCI (2.79 g, 14.5 mmol) and stirred for 10 minutes, followed by addition of HCl-H<sub>2</sub>N-Gly-OBn (2.56 g, 13.0 mmol). The resulting solution was stirred at -20°C for 2 hrs, kept refrigerated overnight and then concentrated to dryness, followed by dilution with EtOAc (150 mL). The EtOAc solution was then washed twice with saturated NaHCO<sub>3</sub>, H<sub>2</sub>O, 5% H<sub>3</sub>PO<sub>4</sub>, brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to dryness to give the Compound (1.09) (4.5 g, 94%). LRMS *m/z* MH<sup>+</sup> = 395.1.


Step B:Compound (1.1)



A solution of compound (1.09) (7.00 g, 17.8 mmol) in absolute ethanol (300 mL) was stirred at room temperature under a hydrogen atmosphere in the presence of Pd-C (300 mg, 10%). The reaction progress was monitored by tlc. After 2 h, the mixture was filtered through a celite pad and the resulting solution was concentrated in vacuo to give Compound (1.1) (5.40 g, quantitative). LRMS  $m/z$   $MH^+ = 305.1$ .


### Preparative Example 2

#### 10 Step A      Compound (1.3)




15

A mixture of Compound (1.1) from Preparative Example 1, Step B (1 eq.), Compound (1.2) (Novabiochem, No. 04-12-5147) (1.03 eq.), HOObt (1.03 eq.), N-methylmorpholine (2.2 eq.), and dimethylformamide (70 mL/g) was stirred at -20°C. EDCI (1.04 eq.) was added and the reaction stirred for 48 hr. The reaction mixture was poured into 5% aqueous  $KH_2PO_4$  and extracted with ethyl acetate (2 x). The combined organics were washed with cold 5% aqueous  $K_2CO_3$ , then 5% aqueous  $KH_2PO_4$ , then brine, and the organic layer was dried over anhydrous  $MgSO_4$ . The mixture was filtered, then evaporated and the filtrate dried under vacuum, the residue was triturated with  $Et_2O$ -hexane, and filtered to obtain the title compound (1.3) (86% yield),  $C_{25}H_{39}N_3O_7$  (493.60), mass spec. (FAB)  $M+1 = 494.3$ .

**Step B      Compound (1.4)**


5                    Compound (1.3) from Preparative Example 2, Step A (3.0 g) was treated with 4 N HCl/dioxane (36 mL) and stirred at room temperature for 7 min. The mixture was poured into 1.5 L cold (5°C) hexane and stirred, then allowed to stand at 0°C for 0.5 hr. The mixture was suction-filtered in a dry  
 10 atmosphere, and the collected solid was further dried to afford the title compound (1.4) (2.3 g, 88% yield), C<sub>20</sub>H<sub>31</sub>N<sub>3</sub>O<sub>5</sub>·HCl, <sup>1</sup>H NMR (DMSO-d<sub>6</sub>/NaOD) δ 7.38 (m, 5H), 5.25 (m, 1H), 4.3-4.1 (m, 1H), 3.8 (m, 2H), 3.4-3.3 (m, obscured by H<sub>2</sub>O), 1.7-1.1 (m, 4H), 1.35 (s, 9H), 0.83 (m, 3H).

**15      Preparative Example 3****Compound (1.5)**

20                    Compound (1.3) from Preparative Example 2, Step A, was treated in essentially the same manner as in Preparative Example 7, Step A below to afford Compound (1.5).

**25      Preparative Example 4****Compound (1.6)**

50




Compound (1.5) from Preparative Example 3, was treated in essentially the same manner as in Preparative Example 2, Step B, to afford

5 Compound (1.6).

### Preparative Example 5


Step A                      Compound (2.09)

5



To a solution of dimethylamine hydrochloride (1.61 g, 19.7 mmol), *N*-Boc-phenylglycine, compound **2.08** (4.50 g, 17.9 mmol, Bachem Co. # A-2225), HOObt (3.07 g, 18.8 mmol) and EDCI (4.12 g, 21.5 mmol) in anhydrous DMF (200 mL) and CH<sub>2</sub>Cl<sub>2</sub> (150 mL) at -20°C was added NMM (5.90 mL, 53.7 mmol). After being stirred at this temperature for 30 min, the reaction mixture was kept in a freezer overnight (18 h). It was then allowed to warm to rt, and EtOAc (450 mL), brine (100 mL) and 5% H<sub>3</sub>PO<sub>4</sub> (100 mL) were added. After the layers were separated, the organic layer was washed with 5% H<sub>3</sub>PO<sub>4</sub> (100 mL), saturated aqueous sodium bicarbonate solution (2 X 150 mL), water (150 mL), and brine (150 mL), dried (MgSO<sub>4</sub>), filtered and concentrated in vacuo to afford Compound (2.09) (4.86 g) as a white solid, which was used without further purification.


Step B                      Compound (2.1)




Compound (2.09) from Preparative Example 5, Step A (4.70 g, crude) was dissolved in 4 N HCl (60 mL, 240 mmol) and the resulting solution was stirred at room temperature. The progress of the reaction was monitored by 10 TLC. After 4 h, the solution was concentrated in vacuo to yield Compound (2.1) as a white solid which was used in the next reaction without further purification. LRMS  $m/z$   $MH^+$  = 179.0.

### Preparative Example 6


Step A                      Compound (2.2)



In essentially the same manner as Preparative Example 2, Step A, substituting phenylglycine N,N-dimethylamide hydrochloride in place of phenylglycine *t*-butyl ester hydrochloride, Compound (2.2) was prepared mass spec. (FAB) M+1 = 465.3.


Step B      Compound (2.3)

Compound (2.2) from Step A (1.85 g) was reacted with 4 N HCl/dioxane (50 mL) at room temperature for 1 hr. The mixture was evaporated under vacuum in a 20°C water bath, triturated under isopropyl ether, filtered, and dried to afford Compound (2.3) (1.57 g, 98% yield),  
 10  $C_{18}H_{28}N_4O_4 \cdot HCl$ , mass spec. (FAB)  $M+1 = 365.3$

Preparative Example 715      Step A      Compound (2.4)

20 A solution of Compound (2.2) from Preparative Example 5, Step A (2.0 g) in dichloromethane (60 mL) was treated with dimethylsulfoxide (3.0 mL) and 2,2-dichloroacetic acid (0.70 mL). The stirred mixture was cooled to 5°C and then added 1 M dicyclohexylcarbodiimide/dichloromethane solution (8.5 mL). The cold bath was removed and the mixture stirred for 22 hr. Then  
 25 added 2-propanol (0.5 mL), and stirred for an additional 1 hr. The mixture was

filtered then washed with ice-cold 0.1 N NaOH (50mL), then ice-cold 0.1 N HCl (50 mL), then 5% aqueous KH<sub>2</sub>PO<sub>4</sub>, then saturated brine. The organic solution was dried over anhydrous magnesium sulfate, then filtered. The filtrate was evaporated, and chromatographed on silica gel, eluting with ethyl acetate to afford Compound (2.3) (1.87 g, 94% yield), C<sub>23</sub>H<sub>34</sub>N<sub>4</sub>O<sub>6</sub>, mass spec. (FAB) M+1 = 463.3.

Step B      Compound (2.5)

5


In essentially the same manner as Preparative Example 2, Step B Compound (2.5) was prepared.

Preparative Example 8

10

Step A      Compound (3.3)

## Step A-1. Compound 3.02




15

A solution of Compound 3.01 (4.6 g, prepared from N-Boc-S-methylcysteine, Bachem Biosciences Inc., according to the procedure of Boger, *J. Org. Chem.*, **1988**, 53(3), 487.) in DMF (150 mL) was treated with Cs<sub>2</sub>CO<sub>3</sub> (6.1 g) followed by benzyl bromide (2.3 mL), and the mixture was stirred at room temperature for 4 hr. The mixture was concentrated *in vacuo*, and the residue suspended in EtOAc (200 mL). Then mixture was washed with aqueous 5% KH<sub>2</sub>PO<sub>4</sub>, then with brine, and the organic extract was dried over anhydrous MgSO<sub>4</sub>. The mixture was filtered, and the filtrate evaporated to leave the product 3.02 (6.2 g); [ $\alpha$ ]<sub>D</sub>-33.7 (c 1.3, CHCl<sub>3</sub>).

20




## Step A-2. Compound 3.03



Following the procedure of U.Larsson, et al., *Acta Chem. Scan.*, **1994**,  
 5 48(6), 517-525, a solution of oxone<sup>(R)</sup> (16.4 g, Aldrich Chemical Co.) in water  
 (90 mL) was added slowly to a 0°C solution of Compound 3.02 (6.1 g) in  
 MeOH (150 mL). The mixture was stirred at room temperature for 4 hr., then  
 concentrated to 1/2 volume on a rotary evaporator, cold water (100 mL) was  
 added, and the mixture extracted with EtOAc. The extract was washed with  
 10 brine, and dried over anhydrous MgSO<sub>4</sub>. The mixture was filtered, and the  
 filtrate evaporated to leave the product 3.03 (5.9 g); [ $\alpha$ ]<sub>D</sub> -26.3 (c 0.9, CHCl<sub>3</sub>).


## Step A-3. Compound 3.2



15 The product 3.03 of the preceding step was treated with 4N  
 HCl/dioxane for 0.5 hr. to afford the product 3.2, C<sub>12</sub>H<sub>17</sub>NO<sub>4</sub>S·HCl (307.79);  
 mass spec. (FAB) M+1 = 272.0.

## Preparation of Compound 3.3


57



Compound (3.2) (S-methyl cysteine sulfone benzyl ester hydrochloride) and Compound (3.1) (N-Boc-cyclohexylglycine) were reacted in essentially the same manner as Preparative Example 2, Step A, to afford Compound (3.3) C<sub>25</sub>H<sub>38</sub>N<sub>2</sub>O<sub>7</sub>S (510.64).


Step B                      Compound (3.4)

10



A mixture of Compound (3.3) from Step A above (0.7 g), 10% Pd/C (0.05 g), and EtOH-dioxane (100 mL) was stirred under 3 atm. H<sub>2</sub> for 5 hr. The mixture was filtered and evaporated to dryness under vacuum to afford compound (3.4) (0.56 g, 97% yield), C<sub>18</sub>H<sub>32</sub>N<sub>2</sub>O<sub>7</sub>S (420.52) mass spec. (FAB) M+1 = 421.2.


Step C                      Compound (3.5)



Compound (3.4) from Step B, above, was reacted with Compound (1.4) from Preparative Example 2, Step B in essentially the same manner as Preparative Example 2, Step A to afford Compound (3.5),  $C_{38}H_{61}N_5O_{11}S$  (795.98), mass spec. (FAB)  $M+1 = 796.3$ .

### Preparative Example 9


### Compound 4.1




Compound (3.5) from Preparative Example 8, Step C, is reacted in essentially the same manner as Preparative Example 2, Step B, to obtain Compound (4.1)  $C_{33}H_{59}N_5O_9S.HCl$  (732.33).

### **Preparative Example 10**


## Compound (4.2)



Treat a solution of Compound (4.1) from Preparative Example 9 (0.7 g), dimethylformamide (15 mL), and diisopropylethylamine (0.38 mL) at 5°C with isobutyl chloroformate (0.15 mL). Remove the cold bath and stir for 6 hr. Pour the reaction mixture into 5% aqueous  $\text{KH}_2\text{PO}_4$  (100 mL) and extract with ethyl acetate (2 x 100 mL). Wash the combined organics with cold 5% aqueous  $\text{K}_2\text{CO}_3$ , then 5% aqueous  $\text{KH}_2\text{PO}_4$ , then brine, and dry the organic over anhydrous  $\text{MgSO}_4$ . Filter the mixture, evaporate the filtrate under vacuum, triturate the residue with  $\text{Et}_2\text{O}$ -hexane, and filter to leave Compound (4.2).


Preparative Example 11Compound (4.3)

Compound (4.2) is reacted in essentially the same manner as Preparative Example 14, Step H below to obtain Compound (4.3).

**Preparative Example 12****Compound (4.4)**

10 Compound (4.3) from Preparative Example 11 (about 0.10 g) is treated with a solution of anhydrous trifluoroacetic acid-dichloromethane (1:1, about 10 mL) for about 2 hr. The solution is diluted with xylene (about 50 mL) and evaporated under vacuum. The residue is triturated with Et<sub>2</sub>O, and filtered to leave Compound (4.4).


15 **Preparative Example 13**

**Compound (4.5)**

20 Compound (4.4) from Preparative Example 12 is reacted with dimethylamine, in essentially the same manner as Preparative Example 2, Step A, to afford Compound (4.5).

**Preparative Example 14****Step A      Compound (5.2)**

5




15 To a stirred cooling solution of Compound (5.01) (1.11 g, 7.0 mmol) in anhydrous DMF (10 mL) and anhydrous  $\text{CH}_2\text{Cl}_2$  (10 mL) at  $0^\circ\text{C}$ , was added HOBT (1.19 g, 7.25 mmol), N-methyl morpholine (2.3 mL, 21.0 mmol) and DEC (1.6 g, 8.4 mmol). The resulting solution was stirred at  $0^\circ\text{C}$  for 15 minutes, followed by the addition of H-Val-O-<sup>t</sup>Bu (1.54 g, 7.35 mmol). The solution was kept in the freezer overnight. A lot of precipitates were observed and the solution was concentrated to dryness, followed by extraction with EtOAc-saturated  $\text{NaHCO}_3$ . The combined organic layer was then washed with 5%  $\text{H}_3\text{PO}_4$  solution,  $\text{H}_2\text{O}$ , brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered to give a crude product 5.1 (2.4 g, 98% yield).

20

The solution of the crude product (as obtained above) in 4N HCl/Dioxane was stirred at room temperature for 7 hrs and concentrated to dryness to afford Compound (5.2).


25

Step B                      Compound (5.4)



To a stirred solution of Compound (5.3) (F. L. Bach, Jr. *et al*, *J. Amer. Chem. Soc.*, (1955) 77, 6049) (17.5 g, 0.086 mmol) in 50%MeOH/50%H<sub>2</sub>O (300 mL) was added Boc anhydride (47.0 g, 0.215 mol). The solution was then adjusted to pH=9.5 by dropwise addition of 50% concentrated NaOH solution. The resulting solution was stirred overnight at room temperature and then neutralized with concentrated HCl to pH=8 and acidified with citric acid to pH=2.94, followed by extraction with CH<sub>2</sub>Cl<sub>2</sub>. The combined organic layer was dried over MgSO<sub>4</sub> to give Compound (5.4) (27.16 g, 95% yield)


Step C                      Compound (5.5)



A solution of thionyl chloride (3.37 mL, 0.046 mmol) in DMF (3.59 mL, 0.046 mol) at 0°C was warmed to room temperature and stirred for 35 minutes. The solution was then cooled to 0°C, followed by addition of  
20 Compound (5.4) from Step B above (15.0 g, 0.045 mol) in CH<sub>3</sub>CN (150 mL) and pyridine (3.73 mL, 0.046 mol). The resulting solution was warmed to room temperature and stirred overnight. The solution was then poured into ice

water (700 mL) and extracted three times with EtOAc (150 mL). The combined organic layer was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to dryness to give Compound (5.5) (10.8 g).


5     Step D             Compound (1.08)



To a stirred solution of Compound (6.6) from Preparative Example 14, Step D below (6.5 g, 0.044 mol) in  $\text{CH}_2\text{Cl}_2$  (130 mL) was added Boc anhydride (9.65 g, 0.044 mol) and DMF (50 mL). The resulting solution was stirred at room temperature over the weekend and concentrated to dryness, followed by addition of  $\text{H}_2\text{O}$  (120 mL) and 50% NaOH to adjust to pH= 10-11. The solution was then stirred for 2 hrs and more Boc anhydride (1.93 g, 8.8 mmol) was added and stirred overnight at room temperature. The solution was then extracted with  $\text{CH}_2\text{Cl}_2$  and the aqueous layer was acidified to pH=4 at  $0^\circ\text{C}$  with 1N HCl and then extracted 3 times with  $\text{CH}_2\text{Cl}_2$ . The combined organic layer was then washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to dryness to give Compound (1.08) ( 4.50 g, 41% yield, M-t-Butyl +2=192).




**Step E**                      **Compound (5.7)**



5

To a stirred solution of Compound (1.08) from above (4.5 g, 0.018 mol) in DMF (22 mL) and CH<sub>2</sub>Cl<sub>2</sub> (22 mL) was added HOBT (2.7 g, 0.02 mol), N-methyl morpholine (6 mL, 0.054 mol), DEC (4.17 g, 0.022 mol) and allyl glycine. TsOH (6.1 g, 0.02 mol). The resulting solution was stirred at room temperature over the weekend and then concentrated to dryness, followed by extraction with EtOAc-saturated NaHCO<sub>3</sub>. The combined organic layer was washed with 10% H<sub>3</sub>PO<sub>4</sub>, brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and filtered to give a crude product (5.7 g). The solution of this crude product in 4N HCl/Dioxane (50 mL) was stirred at room temperature 50 minutes and concentrated to dryness to give Compound (5.7) (4.79 g, 94% yield, MH<sup>+</sup>=245.1).


Step F.            Compound (5.8)



20


To a stirred solution of Compound (5.7) from Step E above (3.1 g, 0.011 mol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (55 mL) was added dropwise TEA (1.69 mL, 0.012 mol) over 13 minutes and Compound (5.5) from Step C (2.83 g, 0.011 mol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (55 mL). The resulting solution was stirred at room temperature for 1.5 hrs. The organic layer was then washed with saturated  $\text{NaHCO}_3$ , brine, dried over  $\text{Na}_2\text{SO}_4$ , and filtered to give Compound (5.8) (4.67 g,  $\text{MH}^+=457.2$ )

Step G      Compound (5.9)



To a stirred solution of Compound (5.2) from step A (0.34 g, 1.31 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 mL) and DMF (5 mL) at  $0^\circ\text{C}$  was added HOBT (0.214 g, 1.31 mmol), N-methyl morpholine (0.43 mL, 3.9 mmol) and DEC (0.5 g, 1.09 mmol). The mixture stirred at room temperature for 15 minutes, followed by the addition of Compound (5.8) from Step F (0.5 g, 1.09 mmol). The resulting solution was kept in the freezer overnight then concentrated to dryness, followed by extraction with EtOAc- $\text{H}_2\text{O}$ . The combined organic layer was washed twice with saturated  $\text{NaHCO}_3$ , 5%  $\text{H}_3\text{PO}_4$ , and brine, dried over  $\text{Na}_2\text{SO}_4$ , then filtered and concentrated to dryness to give Compound (5.9) (0.65 g,  $\text{MH}^+=697.4$ )


Step H                      Compound (5.10)



To a stirred solution of Compound (5.9) from Step G (0.6 g, 0.8 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (8 mL) was added Dess-Martin reagent (0.732 g, 1.72 mmol) and stirred at room temperature for 1 hour, followed by dropwise addition of  $\text{H}_2\text{O}$  (0.031 mL) and Dess-Martin reagent (0.373 g, 0.86 mmol) in  $\text{CH}_2\text{Cl}_2$  (12 mL). The resulting solution was stirred at room temperature for 2.5 hrs, followed by the addition of a solution of 50% saturated  $\text{NaHCO}_3$ /50% saturated  $\text{Na}_2\text{S}_2\text{O}_3$  (20 mL) and stirred rapidly for 1.5 hrs at room temperature. The solution was then washed with  $\text{H}_2\text{O}$ , and brine, dried over  $\text{Na}_2\text{SO}_4$  and concentrated to dryness to give Compound (5.10) (0.588 g, 100% yield,  $\text{MH}^+=695.2$ ).

### **Preparative Example 15**


20	<u>Step A</u>	<u>Compound (6.2)</u>
----	---------------	-----------------------



To a stirred solution of Compound (6.1) (5.0 g, 19.89 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (20 mL) and DMF (10 mL) at -20°C, was added HOBT (3.25 g, 19.89 mmol),


EDCI (4.58 g, 23.87 mmol), and N-methyl morpholine (6.56 mL, 59.69 mmol). The resulting solution was stirred at room temperature for 10 minutes, followed by the addition of  $\text{NH}_4\text{Cl}$  (1.38 g) and kept at 0°C overnight. The solution was then concentrated, and extracted with EtOAc-H<sub>2</sub>O. The combined organic layer was washed twice with  $\text{NaHCO}_3$ ,  $\text{H}_3\text{PO}_4$ , and brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to dryness. Crude product was purified by column chromatography, eluting with 2.5% MeOH/97.5%  $\text{CH}_2\text{Cl}_2$  to give Compound (6.2) (1.95 g,  $\text{MH}^+=251.1$ ).

**Step B**      **Compound (6.3)**



A solution of Compound (6.2) from Step A (12.32 g, 49.28 mmol) in 4N HCl/ Dioxane (270 mL, 43.08 mmol) was stirred at room temperature for 2 hrs then concentrated to dryness to give Compound (6.3) (8.40 g, 100% yield).


**Step C**      **Compound (1.08) (alternate synthesis)**



To a stirred solution of 1-nitrobutane (16.5 g, 0.16 mol) and glyoxylic acid. H<sub>2</sub>O (28.1 g, 0.305 mol) in MeOH ( 122 mL) at 0°C-5°C, was added dropwise triethyl amine (93 mL, 0.667 mol) over 2 hrs. The solution was warmed to room temperature , stirred overnight then concentrated to dryness


to give an oil. The oil was then mixed with H<sub>2</sub>O and acidified to pH-1 with 10% HCl, followed by extraction with EtOAc. The combined organic layer was washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to dryness to give Compound (6.5) ( 28.1 g, 99% yield).

5    Step D            Compound (6.6)



To a stirred solution of Compound (6.5) from Step C (240 g, 1.35 mmol) in acetic acid (1.25 L) was added 10% Pd/C (37 g). The resulting  
10    solution was hydrogenated at 59 psi for 3 hrs and then at 60 psi for overnight. The acetic acid was then evaporated and the residue was azeotroped 3 times with toluene, then triturated with MeOH and ether. The solution was then filtered and azeotroped twice with toluene to give Compound (6.6) (131 g, 66% yield).


15    Step E            Compound (1.08)



To a stirred solution of Compound (6.6) from Step D (2.0 g, 0.0136 mol) in dioxane (10 mL) and H<sub>2</sub>O (5mL) at 0°C, was added 1N NaOH  
20    (aqueous) solution (4.3 mL, 0.014 mol). The resulting solution was stirred for 10 minutes, followed by the addition of Boc anhydride (0.11 g, 0.014 mol) and then stirred at 0°C for 15 minutes. The solution was warmed to room temperature, stirred for 45 minutes, kept in the refrigerator overnight and then concentrated to dryness to give a crude material. To the solution of this crude

material in EtOAc and Ice, was added  $\text{KHSO}_4$  (3.36 g) and  $\text{H}_2\text{O}$  (32 mL) and stirred for 4-6 minutes. The organic layer was then separated and the aqueous layer was extracted twice with EtOAc. The combined organic layer was washed with  $\text{H}_2\text{O}$ , brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to dryness to give Compound (1.08) (3.0 g, 89.2% yield).

Step F      Compound (1.09)




To a stirred solution of Compound (1.08) from Step E (3.0 g, 0.012 mol) in DMF (15 mL) and  $\text{CH}_2\text{Cl}_2$  (15 mL) at  $-20^\circ\text{C}$ , was added HOBT (1.97 g, 0.012 mol), N-methyl morpholine (4.0 mL, 0.036 mol) and EDCI (2.79 g, 0.0145 mol). The reaction stirred for 10 minutes, followed by the addition of H-Gly-OBZ.HCl (2.56 g, 0.013 mol). The resulting solution was stirred at  $-20^\circ\text{C}$  for 2 hrs, kept in the refrigerator overnight, then concentrated to dryness, followed by dilution with EtOAc. The EtOAc solution was then washed twice with saturated  $\text{NaHCO}_3$ ,  $\text{H}_2\text{O}$ , 5%  $\text{H}_3\text{PO}_4$ , brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to dryness to give Compound (1.09) (4.5 g, 94% yield,  $\text{MH}^+ = 395.1$ ).

15


20

Step G                      Compound (6.9)



A solution of Compound (1.09) from Step F (4.5 g, 0.0114mol) in 4N HCl/Dioxane (45 mL) was stirred at room temperature for 45 minutes then concentrated to dryness to give Compound (6.9) (4.5 g,  $MH^+ = 295.1$ ).

Step H                      Compound (6.11)




To a stirred solution of Boc-phenyl-glycine Compound (6.1) ( 0.398 g, 1.58 mmol) in CH<sub>2</sub>Cl<sub>2</sub> (5 mL) and DMF (5 mL) in 100 mL round bottom flask at -20°C, was added HOBt (0.258 g, 1.58 mmol), EDCI (0.364 g, 1.903 mmol), and N-methyl morpholine ( 0.523 mL, 4.759 mmol). The mixture stirred for 5 minutes, followed by the addition of Compound (6.9) from step G (0.5 g, 1.51 mmol) and CH<sub>2</sub>Cl<sub>2</sub> (5 mL). The resulting solution was stirred at -20°C for 10 minutes and then kept in the freezer overnight. The reaction was concentrated to dryness, followed by extraction with EtOAc-saturated NaHCO<sub>3</sub>. The

combined organic layer was then washed twice with 5%  $\text{H}_3\text{PO}_4$ , and brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to dryness to give Compound (6.11) (0.75 g, 94% yield,  $\text{MH}^+=528.1$ ).

Step J      Compound (6.12)


5



A solution of Compound (6.11) from Step H (0.75 g, 1.423 mmol) in 4N HCl/Dioxane (21 mL) was stirred at room temperature for 3 hrs then concentrated to dryness to give Compound (6.12) (0.68 g, 100% yield).

Step K      Compound (6.14)

15




To a stirred solution of Compound (6.13) (0.44 g, 1.725 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 mL) and DMF (5 mL) at  $-20^\circ\text{C}$  was added EDCI (0.39 g, 2.07 mmol), HOBT (0.18 g, 1.725 mmol), and N-methyl morpholine (0.523 mL, 4.76 mmol). The reaction stirred for 5 minutes, followed by the addition of Compound (6.12) from Step J (0.68 g, 1.64 mmol) in  $\text{CH}_2\text{Cl}_2$  (7 mL). The resulting solution was stirred at  $-20^\circ\text{C}$  for 10 minutes and kept in the freezer

20



overnight then concentrated to dryness, followed by extraction with EtOAc-sat.  $\text{NaHCO}_3$ . The combined organic layer was washed twice with 5%  $\text{H}_3\text{PO}_4$ , then brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to dryness to give Compound (6.14) (0.59 g, 54% yield,  $\text{MH}^+ = 667.3$ ).


5    Step L            Compound (6.15)



10            To a stirring solution of Compound (6.14) from Step K (0.593 g, 0.89 mmol) in  $\text{CH}_2\text{Cl}_2$  (20 mL) was added Dess-Martin periodinane (0.76 g, 1.784 mmol). The resulting solution was stirred at room temperature for 2 hrs, followed by the addition of a mixture of  $\text{H}_2\text{O}/\text{CH}_2\text{Cl}_2$ . The mixture stirred for 45 minutes then added 50% saturated  $\text{NaHCO}_3/50\%\text{Na}_2\text{S}_2\text{O}_3$  (10 mL) and  
15            stirred for an additional 1.5 hrs. Additional  $\text{CH}_2\text{Cl}_2$  was added to the solution and the organic layer was washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered, concentrated to dryness, and purified by column chromatography, eluting with 2.5% MeOH/97.5%  $\text{CH}_2\text{Cl}_2$  to give Compound (6.15) (0.48 g, 82% yield).

Step M            Compound (6.16)

20




## 10




To a stirred solution of Compound (6.17) (0.5 g, 1.59 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 mL) and DMF (5 mL) at  $-20^\circ\text{C}$  was added HOBt (0.259 g, 1.59 mmol), NMM (0.48 g, 4.77 mmol), and EDCI (0.366 g, 1.91 mmol). The mixture stirred for 5 minutes, followed by the addition of Compound (6.9) from Step G (0.5 g, 1.51 mmol) and  $\text{CH}_2\text{Cl}_2$  (5 mL). The resulting solution was stirred at  $-20^\circ\text{C}$  for 10 minutes, kept in the freezer overnight then concentrated to dryness, followed by extraction with EtOAc-sat.  $\text{NaHCO}_3$ . The combined organic layer was then washed twice with 5%  $\text{H}_3\text{PO}_4$ , and brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to dryness to give Compound (6.18) (0.95 g,  $\text{MH}^+ = 592.1$ ).

Step O	Compound (6.19)
--------	-----------------



The solution of Compound (6.18) from Step N (0.93 g, 1.58 mmol) in 4N HCl/Dioxane (26 mL) was stirred at room temperature for 2 hr, then concentrated to dryness to give Compound (6.19) (0.96 g, 100% yield,  $MH^+=492.1$ ).


Step P                      Compound (6.20)




To a stirred cooling solution of Compound (6.13) (0.51 g, 2.02 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 mL) and DMF (5 mL) at  $-20^\circ\text{C}$ , was added HOBT (0.33 g, 2.02 mmol), N-methyl morpholine (0.61 g, 6.06 mmol), and EDCI (0.46 g, 2.42 mmol). The reaction stirred for 5 minutes, followed by the addition of

Compound (6.19) from Step O (0.94 g, 1.92 mmol). The resulting solution was stirred at -20°C for 10 minutes, then refrigerated overnight, concentrated to dryness, followed by extraction with EtOAc-sat. NaHCO<sub>3</sub>. The combined organic layer was washed with 5% H<sub>3</sub>PO<sub>4</sub>, and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to dryness to give Compound (6.20) (1.29 g, 87% yield).

Step Q                      Compound (6.21)




To a stirred solution of Compound (6.20) from Step P (1.27 g, 1.74 mmol) in absolute EtOH (50 mL), was added Pd/C (100 mg). The resulting solution was stirred vigorously, followed by addition of 2 drops of AcOH. The solution was then hydrogenated for 2 hrs and filtered through celite to give Compound (6.21) ( 1.07 g, 96% yield,  $MH^+ = 641.1$ ).

Step R      Compound (6.22)

To a stirred solution of Compound (6.21) from Step Q (0.25 g, 0.39 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 mL) and DMF (5 mL) at  $-25^\circ\text{C}$ , was added HOBT (0.06 g, 0.39 mmol), N-methyl morpholine (0.12 g, 1.17 mmol), EDCI (0.089 g, 0.469 mmol) and stirred for 10 minutes, followed by the addition of Compound (6.3) from Step B (0.069 g, 0.37 mmol). The resulting solution was stirred at  $-25^\circ\text{C}$  for 15 minutes, refrigerated over night then concentrated to dryness, followed by extraction with EtOAc-sat.  $\text{NaHCO}_3$ . The combined organic layer was washed with 5%  $\text{H}_3\text{PO}_4$ , then brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to dryness to give Compound (6.22).

10


15

Step S      Compound (6.23)

To a stirred solution of Compound (6.22) from Step R (0.23 g, 0.302 mmol) in  $\text{CH}_2\text{Cl}_2$  (10 mL) was added Dess-Martin periodinane (0.256 g, 0.60 mmol). The resulting solution was stirred at room temperature for 2 hrs, followed by the addition of a mixture of  $\text{H}_2\text{O}/\text{CH}_2\text{Cl}_2$  and stirred for an additional 45 minutes. To the reaction was then added 50% sat.  $\text{NaHCO}_3$ /50%  $\text{Na}_2\text{S}_2\text{O}_3$  (10 mL) and it was stirred for 1.5 hrs. Additional  $\text{CH}_2\text{Cl}_2$  was added to the solution and the organic layer was then washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered, concentrated to dryness and purified by column chromatography on silica gel, eluting with 1%-3%  $\text{MeOH}/99\%$ -97%  $\text{CH}_2\text{Cl}_2$  to give Compound (6.23) (0.08 g, 34% yield,  $\text{MH}^+=771.2$ )


### Preparative Example 16

#### Step A      Compound (7.2)




To a stirred solution of Compound (7.1) (0.476 g, 1.51 mmol) in  $\text{CH}_2\text{Cl}_2$  (60 mL) and DMF (60 mL) at  $-20^\circ\text{C}$ , was added HOBT (0.246 g, 1.51 mmol), N-methyl morpholine (0.458 g, 4.53 mmol), and EDCI (0.351 g, 1.81 mmol). The reaction stirred for 5 minutes, followed by the addition of Compound (6.8) from Preparative Example 15, Step F (0.5 g, 1.51 mmol). The resulting solution was stirred at  $-20^\circ\text{C}$  for 3 hrs, then kept in the freezer overnight, concentrated to dryness, followed by extraction with  $\text{EtOAc}$ -sat.  $\text{NaHCO}_3$ . The combined organic layer was washed twice with 5%  $\text{H}_3\text{PO}_4$ ,  $\text{H}_2\text{O}$ , then brine, dried over  $\text{Na}_2\text{SO}_4$  and concentrated to dryness to give Compound (7.2) (0.82 g, 94% yield,  $\text{MH}^+=592.1$ )

Step B                      Compound (7.3)



A solution of Compound (7.2) from Step A (0.82 g, 1.39 mmol) in 4N HCl/Dioxane (20 mL) was stirred at room temperature for 2 hrs and then concentrated to dryness to give Compound (7.3) (0.84 g, 100% yield,  $MH^+=492.3$ )


Step C                      Compound (7.4)



To a stirred solution of Compound (6.13) (0.36 g, 1.40 mmol) in  $\text{CH}_2\text{Cl}_2$  (60 mL) and DMF (60 mL) at  $-20^\circ\text{C}$  was added HOBT (0.228 g, 1.40 mmol), NMM (0.425 g, 4.20 mmol), and EDCI (0.322 g, 1.68 mmol). The reaction stirred for 5 minutes, followed by the addition of Compound (7.3) from Step B (0.84 g, 1.40 mmol). The resulting solution was stirred at  $-20^\circ\text{C}$  for 3 hrs and then kept in the freezer overnight, concentrated to dryness, followed by extraction with EtOAc-sat.  $\text{NaHCO}_3$ . The combined organic layer was washed


with 5%  $\text{H}_3\text{PO}_4$ ,  $\text{H}_2\text{O}$ , brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to dryness to give Compound (7.4) (0.57 g, 57% yield,  $\text{MH}^+=731.3$ ).

5     Step D                      Compound (7.5)



To a stirred solution of Compound (7.4) from Step C (0.55 g, 0.75 mmol) in  $\text{CH}_2\text{Cl}_2$  (5 mL) was added Dess-Martin periodinane (0.64 g, 1.50 mmol). The resulting solution was stirred at room temperature for 2 hrs, followed by the addition of a mixture of  $\text{H}_2\text{O}/\text{CH}_2\text{Cl}_2$ . The mixture stirred for 45 minutes then added 50% sat.  $\text{NaHCO}_3$ /50%  $\text{Na}_2\text{S}_2\text{O}_3$  and stirred for an additional 1.5 hrs. Additional  $\text{CH}_2\text{Cl}_2$  was added to the solution and the organic layer was then washed with brine, dried over  $\text{Na}_2\text{SO}_4$  and concentrated to dryness to give Compound (7.5) (0.24 g, 44% yield,  $\text{MH}^+ = 729.5$ ).


Step E                      Compound (7.6)





To a stirred solution of Compound (7.5) from Step D (0.10 g, 0.14 mmol) in absolute EtOH (20 mL), was added Pd/C (20 mg). The resulting solution was stirred vigorously in 100 ml round bottom flask, purged with H<sub>2</sub> and stirred under H<sub>2</sub> atmosphere over night. The solution was then filtered through celite, washed with EtOH and concentrated to dryness to give Compound (7.6) (93 mg, 100% yield, MH<sup>+</sup>=639.1).

#### Preparative Example 17




15

In essentially the same manner as Preparative Example 16 Steps A-E, substituting Compound (8.1) for Compound (7.1) in Step A, Compounds (8.2) and (8.3) were prepared.

### Preparative Example 18


5      Step A              Compound (9.2)



10


To a stirred solution of (S) (+)-2-phenyl glycine (9.1) (15.0 g, 0.099 mol) in benzene (350 mL) was added p-toluene sulfonic acid. H<sub>2</sub>O (20.76 g, 0.116 mol) and benzyl alcohol (30 mL, 0.29 mol). The resulting solution was heated to reflux overnight and the solution became a slurry. The solution was then cooled to room temperature, followed by the addition of ether. The solid was filtered through a scintered funnel and washed twice with Et<sub>2</sub>O, then dried under a nitrogen atmosphere to give a solid (35.4 g). The solid was then dissolved in CH<sub>2</sub>Cl<sub>2</sub> and washed with sat. NaHCO<sub>3</sub>. The combined organic layer was dried over Na<sub>2</sub>SO<sub>4</sub> and concentrated to dryness to give a free amine (18.1 g, 75.7% yield). The free amine was then dissolved in ether and 1N HCl was bubbled in to form a white precipitate. The precipitate was filtered, washed with ether and dried under vacuum to give Compound (9.2) (15.2 g).

25      Step B                      Compound (9.4)




To a stirred solution of Boc-gly-OH (9.3) (11.35 g, 0.0648 mol) in anhydrous DMF (100 mL) and anhydrous  $\text{CH}_2\text{Cl}_2$  (100 mL) at  $-20^\circ\text{C}$ , was added HOBT (10.5 g, 0.065 mol), EDCI (13.6 g, 0.0712 mol) and N-methyl morpholine (21.3 mL, 0.194 mol). The resulting solution was stirred at  $-20^\circ\text{C}$  for 10 minutes, followed by the addition of Compound (9.2) from Step A (18.0 g, 0.065 mol). The reaction stirred for 45 minutes at  $-20^\circ\text{C}$  and was then kept in the freezer over night. The solution was then concentrated to dryness, followed by extraction with EtOAc-sat.  $\text{NaHCO}_3$ . The combined organic layer was washed with  $\text{H}_2\text{O}$ , then brine, dried over  $\text{Na}_2\text{SO}_4$  and concentrated to dryness to give Compound (9.4) (26.48 g, 100% yield,  $\text{MH}^+=399.2$ ).

Step C      Compound (9.5)




A solution of Compound (9.4) from Step B (26.4 g, 0.065 mol) in 4N HCl/Dioxane (100 mL) was stirred at room temperature for 1 hr and then concentrated to dryness to give Compound (9.5) (22.69 g, 100% yield,  $\text{MH}^+=299.1$ ).

Step D                      Compound (9.6)




To a stirred solution of Compound (1.08) from Preparative Example 15, 10 Step E (15.5 g, 0.0627 mol) in DMF (150 mL) and  $\text{CH}_2\text{Cl}_2$  (150 mL) at  $-20^\circ\text{C}$ , was added HOBT (10.22 g, 0.0626 mol), EDCI (13.2 g, 0.069 mol) and NMM (20.67 g, 0.188 mol). The resulting solution was stirred at  $-20^\circ\text{C}$  for 10 minutes, followed by the addition of Compound (9.5) from Step C (21.0 g, 0.063 mol). The reaction stirred at  $-20^\circ\text{C}$  for 1 hr and then was kept in the 15 freezer overnight. The solution was then concentrated to dryness, followed by extraction with EtOAc-sat.  $\text{NaHCO}_3$ . The combined organic layer was washed with  $\text{H}_2\text{O}$ , 5%  $\text{H}_3\text{PO}_4$ , and brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to dryness to give Compound (9.6) (30.3 g, 92% yield,  $\text{MH}^+=528.1$ ).

**Step E**      **Compound (9.7)**



In essentially the same manner as Preparative Example 18, Step C above, Compound (9.7) was prepared ( 30.0 g, 100% yield,  $MH^+=428.1$ ).


Step F                      Compound (9.9)



To a stirred solution of Boc-His(Z)-OH (9.8) (0.5 g, 1.28 mmol) in DMF (5 mL) and CH<sub>2</sub>Cl<sub>2</sub> (5 mL) at -20°C, was added HOBT (0.209 g, 1.28 mmol), EDCI (0.27 g, 1.41 mmol), and NMM (0.42 mL, 3.85 mmol). The resulting solution was stirred at -20° for 10 minutes, followed by the addition of Compound (9.7) from Step E (0.673 g, 1.28 mmol) and stirred at -20°C for 2hr and then kept in the freezer over night. The solution was then concentrated to dryness, followed by extraction with EtOAc-sat.NaHCO<sub>3</sub>. The combined organic layer was washed with H<sub>2</sub>O, 5% H<sub>3</sub>PO<sub>4</sub>, and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to give Compound (9.9) (0.858 g, 84% yield, MH<sup>+</sup>=799).

Step G                      Compound (9.11)


85



In essentially the same manner as in Preparative Example 18, Step C,

- 5 Compound (9.11) was prepared (0.76 g, 100% yield,  $MH^+ = 699.2$ ).

Step H      Compound (9.12)




10


- To a stirred solution of N-Boc-cyclohexylglycine (0.263 g, 1.026 mmol) in DMF (5 mL) and  $CH_2Cl_2$  (5 mL) at  $-20^\circ C$ , was added HOBT (0.167 g, 1.026 mmol), EDCI (0.216 g, 1.13 mmol) and NMM (0.338 g, 3.078 mmol). The resulting solution was stirred at  $-20^\circ C$  for 10 minutes, followed by the addition of Compound (9.11) from Step G (0.754 g, 1.03 mmol). The reaction stirred at  $-20^\circ C$  for 1 hr and then was kept in the freezer over night. The solution was then concentrated to dryness, followed by extraction with EtOAc-sat.  $NaHCO_3$ .
- 15
- 20 The combined organic layer was then washed with  $H_2O$ , 5%  $H_3PO_4$ , and brine,

dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to dryness to give Compound (9.12) (0.735g,  $\text{MH}^+=938.4$ ).

Step I                      Compound (9.13)




To a stirred solution of the Compound (9.12) from Step H (0.367g, 0.377 mmol) in anhydrous  $\text{CH}_2\text{Cl}_2$  (10 mL), was added Dess-Martin periodinane (0.32 g, 0.75 mmol). The resulting solution was stirred at room temperature for 2 hrs.  $\text{CH}_2\text{Cl}_2$ , sat.  $\text{Na}_2\text{S}_2\text{O}_4$  and sat.  $\text{NaHCO}_3$  were added to the solution and the solution was allowed to stir at room temperature for 1 hr. The organic layer was then separated and washed with  $\text{H}_2\text{O}$ , and brine, dried over  $\text{Na}_2\text{SO}_4$ , filtered and concentrated to dryness to give a crude product (340 mg). The crude product was then purified by column chromatography on silica gel, eluting with  $\text{CH}_2\text{Cl}_2$  and then 4%  $\text{MeOH}/\text{CH}_2\text{Cl}_2$  to give Compound (9.13) (150 mg,  $\text{MH}^+=936.3$ ).

**Step J      Compound (9.14)**

To a stirred solution of Compound (9.13) from Step I (0.15 g, 1.6  
 10 mmol) in absolute EtOH (40 mL) was added 10% Pd/C in 50% H<sub>2</sub>O (w/w).  
 The solution was purged with N<sub>2</sub> and stirred under H<sub>2</sub> balloon for 45 minutes.  
 The catalyst was then filtered through celite, washed with EtOH/CH<sub>2</sub>Cl<sub>2</sub> and  
 then concentrated to dryness to give Compound (9.14) (0.116 g, MH<sup>+</sup>=712.2).

15 **Preparative Example 19**

**Step A      Compound (10.2)**

A suspension of L-3-(1-Napthyl) Alanine (2.0 g, 9.34 mmol) in  
 25 anhydrous EtOH (200 mL) was charged to a 500 ml flask. To the solution  
 was then bubbled in, anhydrous concentrated HCl (2 mL) to dissolve all the  
 solids. The solution was allowed to cool down to room temperature over 45




minutes and then it was concentrated to dryness, followed by the addition of EtOH (50 mL), 10% Pd/C (300 mg) and 5% Rh/C (300 mg). The resulting solution was placed in a parr shaker and hydrogenated at 60 psi. The reaction was then filtered through celite, washed with EtOH and concentrated to dryness to give a crude material (2.4 g,  $MH^+ = 254.2$ ). The crude product was dissolved in  $CH_2Cl_2$  and then washed with sat.  $NaHCO_3$ . The combined organic layer was concentrated to dryness and purified by column chromatography on silica gel, eluting with 5%-20% EtOAc/ $CH_2Cl_2$  to give Compound (10.2) (0.65 g).

10


Step B      Compound (10.3)

15




To a stirred solution of N-Boc-cyclohexylglycine (0.643 g, 2.5 mmol) in DMF (5 mL) and  $CH_2Cl_2$  (5 mL) at  $-20^\circ C$ , was added HOBT (0.407 g, 2.5 mmol), EDCI (0.527 g, 2.75 mmol) and NMM (0.825 mL, 7.5 mmol). The resulting solution was stirred for 10 minutes at  $-20^\circ C$ , followed by the addition of Compound (10.2) from Step A and  $CH_2Cl_2$  (3 mL), and then kept in the freezer over night. The solution was then concentrated to dryness, followed by extraction with EtOAc-sat.  $NaHCO_3$ . The combined organic layer was washed with  $H_2O$ , 5%  $H_3PO_4$ , and brine, dried over  $Na_2SO_4$ , filtered and concentrated to dryness to give Compound (10.3) (1.12 g, 92% yield).

25

Step CCompound (10.4)

To a stirred solution of Compound (10.3) from Step B (1.1 g, 2.25 mmol) in MeOH (30 mL) and H<sub>2</sub>O (7.5 mL) was added LiOH (0.283 g, 6.75 mmol). The resulting solution was stirred at room temperature over night,  
10 followed by the addition of 5% H<sub>3</sub>PO<sub>4</sub>. A precipitate formed and the solution was evaporated to remove most MeOH. Additional CH<sub>2</sub>Cl<sub>2</sub> was added and the CH<sub>2</sub>Cl<sub>2</sub> layer was then separated, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to dryness to give Compound (10.4) (1.068 g, 100% yield, MH<sup>+</sup>=459.1).

15


Step DCompound (10.5)

25

To a stirred solution of Compound (10.4) from Step C (1.0 g, 2.17 mmol) in DMF (10 mL) and CH<sub>2</sub>Cl<sub>2</sub> (10 mL), was added HOBT (0.353 g, 2.17


mmol), EDCI (0.457 g, 2.38 mmol) and NMM (0.715 mL, 6.51 mmol). The resulting solution was stirred at -20°C for 10 minutes, followed by the addition of Compound (9.7) from Preparative Example 18, Step E (1.13 g, 2.17 mmol). The reaction stirred for 0.5 hr. at -20°C and then was kept in the freezer over  
5 night. The solution was then concentrated to dryness, followed by extraction with EtOAc-sat. NaHCO<sub>3</sub>. The combined organic layer was washed with H<sub>2</sub>O, 5% H<sub>3</sub>PO<sub>4</sub>, and brine, dried over Na<sub>2</sub>SO<sub>4</sub>, filtered and concentrated to dryness to give Compound (10.5) (1.8 g, M+Na=890.4).

10 Step E Compound (10.6)



To a stirred solution of Compound (10.5) from Step D (1.8 g, 2.07 mmol) in anhydrous CH<sub>2</sub>Cl<sub>2</sub> (40 mL) was added Dess-Martin periodinane (1.76 g, 4.15 mmol). The resulting solution was stirred at room temperature for 1 hr,  
20 followed by adding dropwise over 1.5 hrs anhydrous CH<sub>2</sub>Cl<sub>2</sub> (40 mL) and H<sub>2</sub>O (0.074 mL) and stirred an additional 2 hrs. To this solution was then added 40 mL of 50% sat. NaHCO<sub>3</sub>/50% sat. Na<sub>2</sub>S<sub>2</sub>O<sub>4</sub> and the resulting solution was stirred vigorously for half an hour. The organic layer was then separated and washed with H<sub>2</sub>O. The combined organic layer was concentrated to dryness  
25 and purified by column chromatography on silica gel, eluting with 2%-3% MeOH/CH<sub>2</sub>Cl<sub>2</sub> to give Compound (10.6) (0.95 g, MH<sup>+</sup>=866.2).

Step F                      Compound (10.7)



In essentially the same manner as in Preparative Example 18, Step K, Compound (10.7) was prepared.

## Examples

Using the procedures of Preparative Example 1, Step A, and Preparative Example 2, Step F, for couplings; Preparative Example 1, Step B, Preparative Example 1, Step F, Preparative Example 2, Step D, and Preparative Example 4, Step J for ester deprotection; Preparative Example 2, Step E, and Preparative Example 4, Step J, for amine deprotection; and Preparative Example 4, Step H, for oxidation of hydroxyamides to ketoamides – together with the  $\alpha$ -amino acids of the above examples or those commercially available or those described in the literature, in the necessary various combinations, the compounds listed in the attached **Table 2** were prepared.

## 20 Solid Phase Synthesis


**General procedure for solid-phase coupling reactions.**

The synthesis was done in a reaction vessel which was constructed from a polypropylene syringe cartridge fitted with a polypropylene frit at the bottom. The Fmoc-protected amino acids were coupled under standard solid-

phase techniques. Each reaction vessel was loaded with 100 mg of the starting Fmoc-Sieber resin (approximately 0.035 mmol). The resin was washed with 2 mL portions of DMF (2 times). The Fmoc protecting group was removed by treatment with 2 mL of a 20 % v/v solution of piperidine in DMF  
5 for 20 min. The resin was washed with 2 mL portions of DMF (4 times). The coupling was done in DMF (2 mL), using 0.12 mmol of Fmoc-amino acid, 0.12 mmol of HATU [ O-(7-azabenzotriazol-1-yl)-1,1,3,3-tetramethyluronium hexafluorophosphate ] and 0.24 mmol of DIPEA (N,N-diisopropylethylamine). After shaking for 2 h, the reaction vessel was drained and the resin was  
10 washed with 2 mL portions of DMF (4 times). The coupling cycle was repeated with the next Fmoc-amino acid or capping group.

**General procedure for solid-phase Dess-Martin oxidation.**

The synthesis was conducted in a reaction vessel which was constructed from a polypropylene syringe cartridge fitted with a polypropylene  
15 frit at the bottom. Resin-bound hydroxy compound (approximately 0.035 mmol) was treated with a solution of 0.14 mmol of Dess-Martin periodinane and 0.14 mmol of t-BuOH in 2 mL of DCM for 4 h. The resin was washed with 2 mL portions of a 20 % v/v solution of iPrOH in DCM, THF, a 50 % v/v solution of THF in water (4 times), THF (4 times) and DCM (4 times).

**Preparative Example 20****N-Fmoc-2',3'-dimethoxyphenylglycine Compound (901)**

5


To a solution of potassium cyanide (1.465 g, 22.5 mmol) and ammonium carbonate (5.045 g, 52.5 mmol) in water (15 mL) was added a solution of 2,3-dimethoxybenzaldehyde **901A** (2.5 g, 15 mmol) in ethanol (15 mL). The reaction mixture was heated at 40 °C for 24 h. The volume of the solution was reduced to 10 mL by evaporating under reduced pressure. Concentrated hydrochloric acid (15 mL) was added and compound **901B** was obtained as a white precipitate. Compound **901B** was isolated by filtration (2.2 g, 9.3 mmol). Compound **901B** was dissolved in 10 % w/w aqueous sodium hydroxide solution (15 mL) and the resulting solution was heated under reflux for 24 h. Concentrated hydrochloric acid was added and the pH was adjusted to neutral (pH 7). The resulting solution containing compound

15

**901C** was evaporated under reduced pressure. The residue was dissolved in 5 % w/w aqueous sodium bicarbonate solution (150 mL). The solution was cooled to 0 °C in an ice bath and 1,4-dioxane (30 mL) and a solution of 9-fluorenylmethyl succinimidyl carbonate (2.7 g, 8 mmol) in 1,4-dioxane (30 mL) was added at 0 °C. The reaction mixture was allowed to warm to room temperature and was stirred at room temperature for 24 h. 1,4-dioxane was evaporated under reduced pressure. The aqueous solution was washed with diethyl ether. Concentrated hydrochloric acid was added and the pH was adjusted to acidic (pH 1). Ethyl acetate was added the organic layer was washed with water and brine. The organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to afford the desired compound **901** as a white foamy solid (3.44 g, 7.9 mmol). MS (LCMS-Electrospray) 434.1 MH<sup>+</sup>.

#### Preparative Example 21

##### Compound (801)



To a solution of N-Fmoc-phenylalanine **801A** (5 g, 12.9 mmol) in anhydrous DCM (22 mL) cooled to -30°C in a dry ice-acetone bath was added N-methylpyrrolidine (1.96 mL, 16.1 mmol) and methyl chloroformate (1.2 mL, 15.5 mmol) sequentially. The reaction mixture was stirred at -30°C for 1 h

and a solution of N,O-dimethylhydroxylamine hydrochloride (1.51 g, 15.5 mol) and N-methylpyrrolidine (1.96 mL, 16.1 mmol) in anhydrous DCM (8 mL) was added. The reaction mixture was allowed to warm to room temperature and was stirred at room temperature overnight. Toluene was added and the  
5 organic layer was washed with dilute hydrochloric acid, aqueous sodium bicarbonate solution and brine. The organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to afford compound **801B** (4 g, 9.29 mmol).

To a solution of Red-Al (6.28 mL, 21.4 mmol) in anhydrous toluene (8  
10 mL) cooled to -20°C in a dry ice-acetone bath was added a solution of compound **801B** (4 g, 9.29 mmol) in anhydrous toluene (12 mL). The reaction mixture was stirred at -20 °C for 1.5 h. The organic layer was washed with dilute hydrochloric acid, aqueous sodium bicarbonate solution and brine. The organic layer was dried over anhydrous sodium sulfate. The solvent was  
15 evaporated under reduced pressure and the crude product **801C** was used in the next reaction without further purification.


To a solution of compound **801C** (approx. 9.29 mmol) in hexane (15 mL) was added a solution of potassium cyanide (24 mg, 0.37 mmol) and tetrabutylammonium iodide (34 mg, 0.092 mmol) in water (4 mL) and acetone  
20 cyanohydrin (1.27 mL, 13.9 mmol) sequentially. The reaction mixture was stirred at room temperature for 24 h. Ethyl acetate was added and the organic layer was washed with water and brine. The organic layer was dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to afford compound **801D** (2.4 g, 6.03 mmol).

25 To a solution of compound **801D** (2.4 g, 6.03 mmol) in 1,4-dioxane (11 mL) was added concentrated hydrochloric acid (11 mL). The reaction mixture was heated at 80 °C for 3 h. Ethyl acetate (25 mL) and water (25 mL) was added. The organic layer was washed with brine and dried over anhydrous sodium sulfate. The solvent was evaporated under reduced pressure to



afford the desired compound **801** as a white foamy solid (2 g, 4.8 mmol). MS (LCMS-Electrospray) 418.1 MH<sup>+</sup>.

**Example 101J** Compound (101J):




**101J**

**Scheme 5**


(SEE NEXT PAGE)

97




Resin-bound compounds **101B**, **101C**, **101D**, **101E**, **101F** and **101G** were prepared according to the general procedure for solid-phase coupling reactions started with 100 mg of Fmoc-Sieber resin (0.035 mmol). Resin-bound compound **101G** was oxidized to resin-bound compound **101H** according to the general procedure for solid-phase Dess-Martin oxidation. The resin-bound compound **101H** was treated with 4 mL of a 2 % v/v solution of TFA in DCM for 5 min. The filtrate was added to 1 mL of AcOH and the solution was concentrated by vacuum centrifugation to provide compound **101J** (0.011 g, 45 % yield). MS (LCMS-Electrospray) 703.2 MH<sup>+</sup>.




Using the solid phase synthesis techniques defined herein above, the following moieties were prepared and used in accordance with the following Formula:





15 -X:




20





5





10



100





5





10

101





5





10

102








5



10

-P3:

103





5



10



104




5







10

-X1:

105




-P2:



One of -P1a and P1b is -H and the other is selected from:


15

106



-P1':

5




-Z-:

10





15

-P2':





20


107






5






10



108



5



Using the above-described procedures, the compounds with their activity data listed in the attached **Table 3** were prepared.

Additionally prepared were compounds as well as their activity data listed in the attached **Table 4**. Their preparation is described below.

General procedures for preparation of compounds in **Table 4** on solid support:

Solid-phase synthesis is useful for the production of small amounts of certain compounds of the present invention. As with the conventional solid-phase synthesis of peptides, reactors for the solid-phase synthesis of peptidyl ketoamides are comprised of a reactor vessel with at least one surface permeable to solvent and dissolved reagents, but not permeable to synthesis resin of the selected mesh size. Such reactors include glass solid phase reaction vessels with a sintered glass frit, polypropylene tubes or columns with frits, or reactor Kans<sup>TM</sup> made by Irori Inc., San Diego CA. The type of reactor chosen depends on volume of solid-phase resin needed, and different reactor types might be used at different stages of a synthesis. The following procedures will be referenced in the subsequent examples:

**Procedure A: Coupling reaction:** To the resin suspended in N-methylpyrrolidine (NMP) (10-15 mL/ gram resin) was added N-Fmoc or N-Boc-amino acid (2 eq), HOAt (2 eq), HATU (2 eq) and diisopropylethylamine (4 eq). The mixture was let to react for 4-48 hours. The reactants were drained and the resin was washed successively with dimethylformamide, dichloromethane, methanol, dichloromethane and diethylether (use 10-15 mL solvent/ gram resin). The resin was then dried *in vacuo*.

**Procedure B: Fmoc deprotection:** The Fmoc-protected resin was treated with 20% piperidine in dimethylformamide (10 mL reagent/ g resin) for 30 minutes. The reagents were drained and the resin was washed successively with dimethylformamide, dichloromethane, methanol, dichloromethane and diethyl ether (10 mL solvent/ gram resin).

**Procedure C: Boc deprotection:** The Boc-protected resin was treated with a 1:1 mixture of dichloromethane and trifluoroacetic acid for 20-60 minutes (10 mL solvent/ gram resin). The reagents were drained and the resin was washed successively with dichloromethane, dimethylformamide, 5% diisopropylethylamine in dimethylformamide, dimethylformamide,  
5 dichloromethane and dimethylformamide (10 mL solvent/ gram resin).


**Procedure D: Semicarbazone hydrolysis:** The resin was suspended in the cleavage cocktail (10 mL/ g resin) consisting of trifluoroacetic acid: pyruvic acid: dichloromethane: water 9:2:2:1 for 2 hours. The reactants were drained  
10 and the procedure was repeated three more times. The resin was washed successively with dichloromethane, water and dichloromethane and dried under vacuum.

**Procedure E: HF cleavage:** The dried peptide-nVal(CO)-G-O-PAM resin (50 mg) was placed in an HF vessel containing a small stir bar. Anisole  
15 (10% of total volume) was added as a scavenger. In the presence of glutamic acid and cysteine amino acids, thioanisole (10%) and 1,2-ethanedithiol (0.2%) were also added. The HF vessel was then hooked up to the HF apparatus (Immuno Dynamics) and the system was flushed with nitrogen for five minutes. It was then cooled down to -78 °C with a dry ice/ isopropanol bath.  
20 After 20 minutes, HF was distilled to the desired volume (10 mL HF/ g resin). The reaction was let to proceed for 1.5 hours at 0°C. Work up consisted of removing all the HF using nitrogen. Dichloromethane was then added to the resin and the mixture was stirred for five minutes. This was followed by the addition of 20% acetic acid in water (4 mL). After stirring for 20 minutes, the  
25 resin was filtered using a fritted funnel and the dichloromethane was removed under reduced pressure. The remaining solution was washed with hexanes (3x) to remove scavengers. Meanwhile, the resin was soaked in 1 mL methanol. The aqueous layer (20% HOAc) was added back to the resin and the mixture was agitated for five minutes and then filtered. The methanol was  
30 removed under reduced pressure and the aqueous layer was lyophilized. The

peptide was then dissolved in 10-25% methanol (containing 0.1% trifluoroacetic acid) and purified by reverse phase HPLC.

**II) Synthesis of Intermediates:**

**Example I. Synthesis of Boc-3-alkylsulfinylalanine:**




To a mixture of sodium hydride (20 mmol, 800 mg of 60% in oil, washed with hexanes) in tetrahydrofuran (30 mL) at 0°C, was added alkylthiol (20 mmol, R= Ph, R= 1-Naphthyl, R= 2-Naphthyl, R= PhCH<sub>2</sub>CH<sub>2</sub> or R= Et) over 10 min. The cooling bath was removed and stirring was continued for 10 min, at which time Boc-2S-aminopropionyl lactone (reference: *Synthetic Communications*, (1995) 25(16), 2475-2482) (3.74 g, 20 mmol) was added. An ice bath was used to keep the temperature from exceeding 30°C. The reaction mixture was stirred at room temperature for 16 h, concentrated and then dissolved in 1M aqueous potassium bisulfate (200 mL) and 1M HCl (40 mL). The mixture was extracted with dichloromethane (2 x 200 mL). The combined organic layers were dried (sodium sulfate), filtered and concentrated. The residue was dissolved in water (200 mL), methanol (30 mL) and potassium carbonate (40 mmol, 5.5 g). Oxone (21 mmol, 13.0 g) was added in portions with cooling to maintain room temperature. The mixture was stirred for 18 hours and then concentrated in vacuo to remove the methanol. The solution was acidified with 2M potassium bisulfate (pH=1) then extracted with ethyl acetate (2 x 100 mL). The combined organic layers were dried (sodium sulfate), filtered and concentrated. Purification was performed using reverse-phase HPLC. After purification the product was




further purified by acid base extraction to remove de-bocylated material and then stored as the diisopropylethyl ammonium salt to prevent further decomposition (loss of Boc group).

**Example II. Synthesis of 2-(1-methylethyl)-7-methyl-oct-4-enoic acid:**




The above intermediate was synthesized according to the published procedure (Wuts, P. G. M.; Ritter, A. R.; Pruitt, L. E. *J. Org. Chem.* (1992) 57, 6696-6700).

**Example III. Synthesis of Fmoc-nV-(dpse)-Gly-OH (Steps 1-7 below)**



**Step 1. Synthesis of allyl isocyanoacetate (steps a-b below)**


**a) Synthesis of isocyanoacetic acid potassium salt:**



Ethyl isocyanoacetate (96.6 ml, 0.88 mol) was added dropwise to a chilled solution of ethanol (1.5 L) and potassium hydroxide (59.52 g, 1.0 mol). The reaction was slowly warmed to room temperature. After two hours the precipitated product was collected by filtration and washed with several


portions of chilled ethanol. The potassium salt of isocyanoacetic acid thus obtained was dried in vacuo to a golden-brown solid (99.92 g, 91.8%).

b) Synthesis of allyl isocyanoacetate:




To the product of part a (99.92 g, 0.81 mol) dissolved in acetonitrile (810 ml) was added allyl bromide (92 ml, 1.05 mol). After heating at reflux for four hours a dark brown solution was obtained. The reaction mixture was concentrated and the remaining residue was dissolved in ether (1.5 L) and washed with water (3 x 500 ml). The organic layer was dried, filtered and concentrated to a dark brown syrup. The crude was purified by vacuum distillation at 7 mm Hg (98°C) to a clear oil (78.92 g, 78%). NMR ppm (CDCl<sub>3</sub>): 5.9 (m, 1 H), 5.3 (m, 2H), 4.7 (d, 2H), 4.25 (s, 2H).

15 **Step 2. Synthesis of 9-fluorenylmethoxycarbonyl-norvalinal (steps a-c below)**




a) Synthesis of 9-fluorenylmethoxycarbonyl-L-norvaline methyl ester  
(Fmoc-nVal-OMe):



To a chilled solution of the commercially available Fmoc-L-norvaline (25 g, 73.75 mmol) in anhydrous methanol (469 ml) was added thionyl chloride (53.76 ml, 737.5 mmol) over one hour. TLC in ethylacetate taken an hour later confirmed the completion of the reaction ( $R_f = 0.85$ ). The reaction mixture was concentrated and the remaining residue was dissolved in ethylacetate. The organic layer was washed with saturated sodium bicarbonate (3 x 200 ml) followed by brine (200 ml). The organic layer was dried, filtered and concentrated to afford Fmoc-norVal-OMe as a white solid (26.03 g) in quantitative yield. NMR ppm ( $CD_3OD$ ): 7.7 (m, 2H), 7.6 (m, 2H), 7.4 (m, 2H), 7.3 (m, 2H), 4.3 (m, 2H), 4.1 (m, 2H), 3.7 (s, 3H), 1.7 (m, 1H), 1.6 (m, 1H), 1.4 (m, 2H), 0.95 (t, 3H).


b) Synthesis of 9-fluorenylmethoxycarbonyl-norvalinol (Fmoc-nValinol):



To Fmoc-nVal-OMe (26.03 g, 73.75 mmol) in tetrahydrofuran (123 ml) and methanol (246 ml) was added calcium chloride (16.37 g, 147.49 mmol). The reaction mixture was cooled to 0°C and sodium borohydride (11.16 g, 294.98 mmol) was added in several batches. To the thick paste obtained 500 ml methanol was added and the reaction was let to stir at room temperature for 90 minutes. TLC in 2:3 ethylacetate: hexane confirmed the completion of the reaction ( $R_f = 0.25$ ). The reaction was quenched with the slow addition of 100 ml water at 0 °C. The methanol was removed under reduced pressure and the remaining aqueous phase was diluted with ethylacetate. The organic layer was washed with water (3 x 500 ml), saturated sodium bicarbonate (3 x 500 ml) and brine (500 ml). The organic layer was dried over sodium sulfate and concentrated to a white solid (21.70 g, 90.5%). NMR ppm ( $CD_3OD$ ):

7.8 (m, 2H), 7.7 (m, 2H), 7.4 (m, 2H), 7.3 (m, 2H), 4.3-4.5 (m, 2H), 4.2 (m, 1H), 3.6 (s, 1H), 3.5 (s, 2H), 1.5 (m, 1H), 1.3-1.4 (m, 3H), 0.99 (m, 3H).

c) Synthesis of 9-fluorenylmethoxycarbonyl-norvalinal (Fmoc-nVal-CHO):



5


To a solution of Fmoc-norValinol (21.70 g, 66.77 mmol) in dichloromethane (668 ml) was added triethylamine (37.23 ml, 267 mmol) and the solution was cooled to 0°C. A suspension of pyridine sulfur trioxide complex (42.51 g, 267 mmol) in dimethylsulfoxide (96 ml) was added to the chilled solution. After one hour, TLC in 2:3 ethylacetate: hexanes confirmed the completion of the reaction. The dichloromethane was removed under reduced pressure and the remaining residue was dissolved in ethylacetate and washed with water (2 × 50 ml), 1N saturated sodium bisulfate (2 × 50 ml), saturated sodium bicarbonate (2 × 50 ml) and brine (50 ml). The organic layer was concentrated to yield a white solid. Theoretical yield (21.57 g) was assumed and the reaction was taken to the next step without further purification.

15

**Step 3. Synthesis of diphenylmethyl semicarbazide (dpssc) trifluoroacetate salt (steps a-b below)**

20

a) Synthesis of Boc-semicarbazid-4-yl diphenylmethane




To a solution of carbonyldiimidazole (16.2 g, 0.10 mole) in dimethylformamide (225 ml) was added a solution of tert-butyl carbazate (13.2 g, 0.100 mol) in dimethylformamide (225 ml) dropwise over 30 minutes. Diphenylmethanimine (18.3 g, 0.10 mol) was added next over 30 minutes.

5 The reaction was allowed to stir at room temperature for one hour. Water (10 mL) was added and the mixture was concentrated to about 150 mL under reduced pressure. This solution was poured into water (500 mL) and extracted with ethyl acetate (400 mL). The ethylacetate phase was washed with 1N HCl (2 x 75 mL), H<sub>2</sub>O (2 x 75 mL), saturated sodium bicarbonate

10 solution (2 x 75 mL) and sodium chloride (2 x 75 mL), and dried with magnesium sulfate. The mixture was filtered and the solution was concentrated to give 29.5 g (85% yield) of a white foam. This material could be purified by recrystallization from ethyl acetate/hexane, but was pure enough to use directly in the next step: mp 142-143°C. <sup>1</sup>H NMR (CDCl<sub>3</sub>) δ

15 1.45 (s, 9H), 6.10 (dd, 2H), 6.42 (s, 1H), 6.67 (bs, 1H), 7.21-7.31 (m, 10H). Analytical calculated for C<sub>19</sub>H<sub>23</sub>N<sub>3</sub>O<sub>3</sub>: C, 66.84; H, 6.79; N, 12.31. Found: C, 66.46; H, 6.75; N, 12.90.

b) Synthesis of diphenylmethyl semicarbazide (dpSC) trifluoroacetate salt




20

A solution of Boc-semicarbazid-4-yl diphenylmethane (3.43 g, 10 mmol) in dichloromethane (12.5 ml) was treated with trifluoroacetic acid (12.5 ml) at room temperature and allowed to stir for 30 min. The solution was added dropwise to 75 mL of ether and the resulting solid (2.7 g, 80%) was

25 collected by filtration. mp 182-184°C. <sup>1</sup>H NMR (CD<sub>3</sub>OD) δ 6.05 (s, 1H),

7.21-7.35 (m, 10H).  $^{13}\text{C}$  NMR ( $\text{CD}_3\text{OD}$ )  $\delta$  57.6, 118.3 (q,  $\text{CF}_3$ ), 126.7, 127.9, 141.6, 156.9, 160.9 (q,  $\text{CF}_3\text{CO}_2\text{H}$ ).

**Step 4. Synthesis of Fmoc-nVal-(CHOH)-Gly-Oallyl**




To a solution of Fmoc-nVal-CHO (Example III, Step 2c) (5.47 g, 16.90 mmol) in dichloromethane (170 ml) was added allyl isocyanoacetate (Example III, Step 1b) (2.46 ml, 20.28 mmol) and pyridine (5.47 ml, 67.61 mmol). The reaction mixture was cooled to 0 °C and trifluoroacetic acid (3.38 ml, 33.80 mmol) was added dropwise. The reaction was stirred at 0 °C for 1h, and then at room temperature for 48 hours. TLC taken in ethylacetate confirmed the completion of the reaction. The reaction mixture was concentrated and subjected to flash chromatography using 20% to 70% ethylacetate in hexanes. Fractions containing the desired product were pooled and concentrated to a white foam (6.88 g, 87.3%). TLC in 1:1 ethylacetate/hexanes showed one spot ( $R_f = 0.37$ ). NMR  $\delta$  ppm ( $\text{CD}_3\text{OD}$ ): 7.8 (m, 2H), 7.65 (m, 2H), 7.4 (m, 2H), 7.3 (m, 2H), 5.9 (m, 1H), 5.1-5.4 (m, 2H), 4.55-4.65 (m, 2H), 4.3-4.4 (m, 2H), 4.15-4.25 (m, 1H), 4.01 (s, 1H), 3.9-4.0 (m, 3H), 1.5-1.6 (m, 2H), 1.35-1.45 (m, 3H), 0.9 (m, 3H).

10

15


20

**Step 5. Synthesis of Fmoc-nVal-(CO)-Gly-Oallyl**



To a solution of Fmoc-nVal-(CHOH)-Gly-Oallyl (Step 4) (5.01 g, 10.77 mmol) in dimethylsulfoxide (100 ml) and toluene (100 ml) was added EDC (20.6 g, 107.7 mmol). The reaction mixture was cooled to 0 °C and dichloroacetic acid (4.44 ml, 53.83 mmol) was added dropwise. The reaction was stirred for 15 minutes at 0 °C and 1h at room temperature. Water (70 ml) was added at 0 °C and the toluene was removed under reduced pressure. The remaining residue was diluted with ethylacetate and washed several times with a saturated sodium bicarbonate solution followed by 1N sodium bisulfate and brine. The organic layer was dried over sodium sulfate, filtered and concentrated. The theoretical yield of 4.99 g was assumed and the reaction was taken to the next step without further purification. TLC in 1:1 ethylacetate/hexanes shows one spot ( $R_f = 0.73$ ).

#### Step 6. Synthesis of Fmoc-nVal-(dpSC)-Gly-Oallyl




15

To a solution of Fmoc-nVal-(CO)-Gly-Oallyl (step 5) (4.99 g, 10.75 mmol) in ethanol (130 ml) and water (42 ml) were added diphenylmethyl semicarbazide (dpSC) trifluoroacetate salt (Example III, Step 3b) (7.6 g, 21.5 mmol) and sodium acetate  $\cdot 3\text{H}_2\text{O}$  (1.76 g, 12.9 mmol). The reaction mixture was heated at reflux for 90 minutes. The completion of reaction was confirmed by thin layer chromatography taken in 1:1 ethylacetate: hexane. Ethanol was removed under reduced pressure and the remaining residue was dissolved in ethylacetate and washed with 1N sodium bisulfate (2 x 10 mL), saturated sodium bicarbonate (2 x 10 mL) and brine (2 x 10 mL). The organic layer was dried, filtered and concentrated. The resulting residue was purified by flash chromatography in 20% to 50% ethylacetate in

25

hexanes to give a white solid (5.76 g, 78%). TLC in 1:1 ethylacetate/hexanes showed two spots (cis and trans isomers) with  $R_f = 0.42$  and 0.50.

**Step 7. Synthesis of Fmoc-nVal-(dpSC)-Gly-OH**





- 5 To a solution of Fmoc-nVal-(dpSC)-Gly-Oallyl (example III; Step 6) (4.53 g, 6.59 mmol) in tetrahydrofuran (300 ml) was added dimedone (4.62 g, 32.97 mmol) followed by tetrakis(triphenylphosphine) palladium(0) catalyst (0.76 g, 0.66 mmol). After 90 minutes the completion of the reaction was confirmed by TLC (9:1 dichloromethane: methanol). The reaction mixture was
- 10 concentrated and the remaining residue was dissolved in ethylacetate and washed with 0.1M potassium biphosphate (3 x 50 mL). The organic layer was then treated with 50 ml sodium bisulfite and the two phase system was stirred for 15 minutes. The phases were separated and the procedure was repeated twice more. The organic layer was dried, filtered and concentrated and
- 15 subjected to flash column chromatography (20% to 100% ethylacetate in hexanes, then 9:1 dichloromethane: methanol) to obtain a white solid (3.99 g, 94%). TLC in 9:1 dichloromethane: methanol shows two spots (cis and trans isomers). NMR  $\delta$  ppm (CD<sub>3</sub>OD): 7.75 (m, 2H), 7.6 (m, 3H), 7.2-7.4 (m, 14H), 6.1-6.2 (m, 1H), 4.25-4.4 (m, 2H), 4.1-4.2 (m, 2H), 3.85 (s, 2H), 1.6-1.8 (m,
- 20 2H), 1.3-1.5 (m, 2H), 0.95 (t, 3H).

**Example IV. Synthesis of H-nVal(dpSC)-Gly-Phg-MBHA resin (Steps 1-2 below)**



120

**Step 1. Synthesis H-Phg-MBHA resin**


5

Commercially available MBHA resin (2.6 g, 1.12 mmol/g, 2.91 mmol) was transferred to a 250 mL fritted solid phase reaction vessel equipped with a nitrogen inlet. It was then washed thoroughly with 30 ml portions of dichloromethane, methanol, dimethylformamide and dichloromethane and coupled over 18 hours to the commercially available Fmoc-Phg-OH (2.17 g, 5.82 mmol) according Procedure A with 99.82% efficiency. The resin was then subjected to Fmoc deprotection according to procedure B. A qualitative ninhydrin assay on a small aliquot gave dark blue resin and solution, indicating a successful reaction.

15

**Step 2. Synthesis of H-nVal(dpse)-Gly-Phg-MBHA resin**

121




The resin obtained in step 1 above (2.6 g, 0.8 mmol/g, 2.91 mmol) was reacted with Fmoc-nVal-(dpse)-Gly-OH Example III, Step 7) (5.82 mmol, 3.77 g) according to Procedure A. After 18 hours, quantitative ninhydrin analysis indicated 99.91% coupling efficiency. The resin was subjected to Fmoc deprotection according to procedure B. A qualitative ninhydrin assay on a small aliquot gave dark blue resin and solution, indicating a successful reaction.

**III) Solid phase assembly of representative Hepatitis C targets:**

**Example V. Solid Phase Synthesis of 2,5-difluoro-6-**


**hydroxycarbonylphenylcarbonyl-G(Chx)-Leu-nVal-(CO)-Gly-Phg-NH<sub>2</sub>**

**(Steps 1-5 below)**



**Step 1. Synthesis of Fmoc-Leu-nVal(dpse)-Gly-Phg-MBHA resin**


122



The compound H-nVal(dpse)-Gly-Phe-MBHA resin (Example IV, Step 2) (1.5 g, 1.12 mmol/g, 1.68 mmol) was transferred to a fritted polypropylene tube and was coupled to N-Fmoc-Leu-OH (890 mg, 2.52 mmol) according to procedure A. After 18 hours, qualitative ninhydrin analysis showed colorless beads and solution.


**Step 2.** Synthesis of Fmoc-G(Chx)-Leu-nVal(dpse)-Gly-Phe-MBHA resin

123



The resin obtained in Example V, Step 1 (Fmoc-Leu-nVal(dpse)-Gly-Phg-MBHA resin, 1.68 mmol) was subjected to the Fmoc deprotection  
 5 procedure according to Procedure B. Commercially available Fmoc-G(Chx)-OH (0.956 g, 0.252 mmol) was then coupled according to procedure A. After 18 hours quantitative ninhydrin analysis indicated 98% coupling efficiency.  
**Step 3.** Synthesis of 2,5-difluoro-6-hydroxycarbonylphenylcarbonyl-G(Chx)-Leu-nVal(dpse)-Gly-Phg-MBHA resin


124



The resin obtained in Example V, Step 2 (Fmoc-G(Chx)-Leu-nVal(dpse)-Gly-Phg-MBHA resin) was subjected to Fmoc deprotection according to procedure B. A ninhydrin assay on a small aliquot gave dark blue resin and solution, indicating a successful reaction. To the resin (150 mg, 0.168 mmol) suspended in 1 ml NMP was added 3,6-difluorophthalic anhydride (91 mg, 0.42 mmol) followed by diisopropylethylamine (0.146 ml, 84 mmol), and the reaction mixture was shaken for 18 hours at room temperature. Quantitative ninhydrin analysis indicated 97.8% coupling efficiency.


**Step 4.** Synthesis of 2,5-difluoro-6-hydroxycarbonylphenylcarbonyl-G(Chx)-Leu-nVal(CO)-Gly-Phg-MBHA resin

125



The compound of step Example V, Step 3 (2,5-difluoro-6-hydroxycarbonylphenylcarbonyl-G(Chx)-Leu-nVal(dpse)-Gly-Phe-MBHA resin) (200 mg) was subjected to semicarbazone hydrolysis Procedure D.


- 5 **Step 5.** Synthesis of 2,5-difluoro-6-hydroxycarbonylphenylcarbonyl-G(Chx)-Leu-nVal(CO)-Gly-Phe-NH<sub>2</sub>




The resin of Example V, Step 4 (2,5-difluoro-6-hydroxycarbonylphenylcarbonyl-G(Chx)-Leu-nVal(CO)-Gly-Phg-MBHA resin) (100 mg) was subjected to HF cleavage condition (Procedure E) to yield the desired crude product. The material was purified by HPLC using a 2.2 × 25 cm reverse phase column, containing a C-18 resin comprised of 10 micron size gel particles with a 300 angstrom pore size, eluting with a gradient using 20-50% acetonitrile in water. Analytical HPLC using a 4.6 × 250 mm reverse phase column, containing a C-18 resin comprised of 5 micron size gel particles with a 300 angstrom pore size, eluting with 10-60% acetonitrile in water (containing 0.1% trifluoroacetic acid) showed one peak at 17.2 minutes. Low resolution mass spectrum confirmed the desired mass (MH<sup>+</sup> 771.5).

**Example VI. Solid Phase Synthesis of iBoc-G(Chx)-Cys((O<sub>2</sub>)Et)-nVal-(CO)-Gly-Phg-NH<sub>2</sub> (Steps 1-5 below)**

127



**Step 1. Synthesis of Fmoc-Cys((O2)Et)-nVal(dpse)-Gly-Phg-MBHA resin**




5

The compound H-nVal(dpse)-Gly-Phg-MBHA resin (Example IV, Step 2) (0.17 g, 0.8 mmol/g, 0.19 mmol) was transferred to a fritted polypropylene tube and was coupled to Boc-Cys((O2)Et)-OH (Example I) (160 mg, 0.38 mmol) according to procedure A. After 18 hours, quantitative ninhydrin analysis showed the coupling was 99.98% complete.

10




**Step 2. Synthesis of Fmoc-G(Chx)-Cys((O2)Et)-nVal(dpse)-Gly-Phg-MBHA resin**



5 The resin obtained in the previous step (Example VI, Step 1) (Boc-Cys((O2)Et)-nVal(dpse)-Gly-Phe-MBHA resin, 0.19 mmol) was subjected to the Boc deprotection procedure according to Procedure C. Fmoc-G(Chx)-OH (0.170 g, 0.45 mmol) was then coupled according to procedure A. After 18 hours quantitative ninhydrin analysis indicated 99.92% coupling efficiency.

10 **Step 3. Synthesis of iBoc-G(Chx)-Cys(O<sup>2</sup>Et)-nVal(dpse)-Gly-Phg-MBHA resin**


129



The resin obtained in the previous step (Example VI, Step 2) (Fmoc-G(Chx)-Cys((O2)Et)-nVal(dpse)-Gly-Phg-MBHA resin) was subjected to Fmoc deprotection according to procedure B. A ninhydrin assay on a small aliquot gave dark blue resin and solution, indicating a successful reaction. To the resin (170 mg, 0.19 mmol) suspended in 1 ml NMP was added isobutyl chloroformate (0.06 mL, 0.45 mmol) followed by diisopropylethylamine (0.16 ml, 0.90 mmol), and the reaction mixture was shaken for 18 hours at room temperature. Quantitative ninhydrin analysis indicated 99.35% coupling efficiency.

**Step 4.** Synthesis of Iboc-G(Chx)-Cys((O2)Et)-nVal(CO)-Gly-Phg-MBHA resin


130



The compound of the previous step (Example VI, Step 3) iBoc-G(Chx)-Cys(O<sup>2</sup>Et)-nVal(dpse)-Gly-Phg-MBHA resin (170 mg) was subjected to semicarbazone hydrolysis Procedure D.

5 **Step 5.** Synthesis of iBoc-G(Chx)-Cys((O<sup>2</sup>Et)-nVal(CO)-Gly-Phg-NH<sub>2</sub>

131



The resin of the previous step (Example VI, Step 4) (iBoc-G(Chx)-Cys((O2)Et)-nVal(CO)-Gly-Phg-MBHA resin) (170 mg) was subjected to HF cleavage condition (Procedure E) to yield the desired crude product. The material was purified by HPLC using a 2.2 × 25 cm reverse phase column, containing a C-18 resin comprised of 10 micron size gel particles with a 300 angstrom pore size, eluting with a gradient using 20-50% acetonitrile in water. Analytical HPLC using a 4.6 × 250 mm reverse phase column, containing a C-18 resin comprised of 5 micron size gel particles with a 300 angstrom pore size, eluting with 10-60% acetonitrile in water (containing 0.1% trifluoroacetic acid) showed one peak at 16.94 minutes. Low resolution mass spectrum confirmed the desired mass ( $MH^+$  737.5).

#### **Assay for HCV Protease Inhibitory Activity:**

**Spectrophotometric Assay:** Spectrophotometric assay for the HCV serine protease was performed on the inventive compounds by following the procedure described by R. Zhang *et al*, *Analytical Biochemistry*, **270** (1999) 268-275, the disclosure of which is incorporated herein by reference. The assay based on the proteolysis of chromogenic ester substrates is suitable for

the continuous monitoring of HCV NS3 protease activity. The substrates were derived from the P side of the NS5A-NS5B junction sequence (Ac-DTEDV $\text{VX}$ (Nva), where X = A or P) whose C-terminal carboxyl groups were esterified with one of four different chromophoric alcohols (3- or 4-nitrophenol, 7-hydroxy-4-methyl-coumarin, or 4-phenylazophenol). Presented below are the synthesis, characterization and application of these novel spectrophotometric ester substrates to high throughput screening and detailed kinetic evaluation of HCV NS3 protease inhibitors.

Materials and Methods:

Materials: Chemical reagents for assay related buffers were obtained from Sigma Chemical Company (St. Louis, Missouri). Reagents for peptide synthesis were from Aldrich Chemicals, Novabiochem (San Diego, California), Applied Biosystems (Foster City, California) and Perseptive Biosystems (Framingham, Massachusetts). Peptides were synthesized manually or on an automated ABI model 431A synthesizer (from Applied Biosystems). UV/VIS Spectrometer model LAMBDA 12 was from Perkin Elmer (Norwalk, Connecticut) and 96-well UV plates were obtained from Corning (Corning, New York). The prewarming block was from USA Scientific (Ocala, Florida) and the 96-well plate vortexer was from Labline Instruments (Melrose Park, Illinois). A Spectramax Plus microtiter plate reader with monochromator was obtained from Molecular Devices (Sunnyvale, California).

Enzyme Preparation: Recombinant heterodimeric HCV NS3/NS4A protease (strain 1a) was prepared by using the procedures published previously (D. L. Sali *et al*, *Biochemistry*, 37 (1998) 3392-3401). Protein concentrations were determined by the Biorad dye method using recombinant HCV protease standards previously quantified by amino acid analysis. Prior to assay initiation, the enzyme storage buffer (50 mM sodium phosphate pH 8.0, 300 mM NaCl, 10% glycerol, 0.05% lauryl maltoside and 10 mM DTT) was exchanged for the assay buffer (25 mM MOPS pH 6.5, 300 mM NaCl, 10%

glycerol, 0.05% lauryl maltoside, 5  $\mu$ M EDTA and 5  $\mu$ M DTT) utilizing a Biorad Bio-Spin P-6 prepacked column.

Substrate Synthesis and Purification: The synthesis of the substrates was done as reported by R. Zhang *et al.*, (*ibid.*) and was initiated by anchoring  
5 Fmoc-Nva-OH to 2-chlorotrityl chloride resin using a standard protocol (K. Barlos *et al.*, *Int. J. Pept. Protein Res.*, 37 (1991), 513-520). The peptides were subsequently assembled, using Fmoc chemistry, either manually or on an automatic ABI model 431 peptide synthesizer. The N-acetylated and fully  
10 protected peptide fragments were cleaved from the resin either by 10% acetic acid (HOAc) and 10% trifluoroethanol (TFE) in dichloromethane (DCM) for 30 min, or by 2% trifluoroacetic acid (TFA) in DCM for 10 min. The combined filtrate and DCM wash was evaporated azeotropically (or repeatedly extracted by aqueous Na<sub>2</sub>CO<sub>3</sub> solution) to remove the acid used in cleavage. The DCM phase was dried over Na<sub>2</sub>SO<sub>4</sub> and evaporated.

15 The ester substrates were assembled using standard acid-alcohol coupling procedures (K. Holmber *et al.*, *Acta Chem. Scand.*, B33 (1979) 410-412). Peptide fragments were dissolved in anhydrous pyridine (30-60 mg/ml) to which 10 molar equivalents of chromophore and a catalytic amount (0.1 eq.) of para-toluenesulfonic acid (pTSA) were added.  
20 Dicyclohexylcarbodiimide (DCC, 3 eq.) was added to initiate the coupling reactions. Product formation was monitored by HPLC and found to be complete following 12-72 hour reaction at room temperature. Pyridine solvent was evaporated under vacuum and further removed by azeotropic evaporation with toluene. The peptide ester was deprotected with 95% TFA in  
25 DCM for two hours and extracted three times with anhydrous ethyl ether to remove excess chromophore. The deprotected substrate was purified by reversed phase HPLC on a C3 or C8 column with a 30% to 60% acetonitrile gradient (using six column volumes). The overall yield following HPLC purification was approximately 20-30%. The molecular mass was confirmed

by electrospray ionization mass spectroscopy. The substrates were stored in dry powder form under desiccation.

Spectra of Substrates and Products: Spectra of substrates and the

corresponding chromophore products were obtained in the pH 6.5 assay buffer. Extinction coefficients were determined at the optimal off-peak wavelength in 1-cm cuvettes (340 nm for 3-Np and HMC, 370 nm for PAP and 400 nm for 4-Np) using multiple dilutions. The optimal off-peak wavelength was defined as that wavelength yielding the maximum fractional difference in absorbance between substrate and product (product OD - substrate

OD)/substrate OD).

Protease Assay: HCV protease assays were performed at 30°C using a 200 µl reaction mix in a 96-well microtiter plate. Assay buffer conditions (25 mM MOPS pH 6.5, 300 mM NaCl, 10% glycerol, 0.05% lauryl maltoside, 5 µM EDTA and 5 µM DTT) were optimized for the NS3/NS4A heterodimer (D. L.

Sall *et al, ibid.*). Typically, 150 µl mixtures of buffer, substrate and inhibitor were placed in wells (final concentration of DMSO 4 % v/v) and allowed to preincubate at 30 °C for approximately 3 minutes. Fifty µls of prewarmed protease (12 nM, 30°C) in assay buffer, was then used to initiate the reaction (final volume 200 µl). The plates were monitored over the length of the assay (60 minutes) for change in absorbance at the appropriate wavelength (340 nm for 3-Np and HMC, 370 nm for PAP, and 400 nm for 4-Np) using a

Spectromax Plus microtiter plate reader equipped with a monochrometer (acceptable results can be obtained with plate readers that utilize cutoff filters). Proteolytic cleavage of the ester linkage between the Nva and the

chromophore was monitored at the appropriate wavelength against a no enzyme blank as a control for non-enzymatic hydrolysis. The evaluation of substrate kinetic parameters was performed over a 30-fold substrate concentration range (~6-200 µM). Initial velocities were determined using linear regression and kinetic constants were obtained by fitting the data to the Michaelis-Menten equation using non-linear regression analysis (Mac Curve

Fit 1.1, K. Raner). Turnover numbers ( $k_{cat}$ ) were calculated assuming the enzyme was fully active.

Evaluation of Inhibitors and Inactivators: The inhibition constants ( $K_i$ ) for the competitive inhibitors Ac-D-(D-Gla)-L-I-(Cha)-C-OH (27), Ac-DTENVVA(Nva)-OH and Ac-DTENVVP(Nva)-OH were determined experimentally at fixed concentrations of enzyme and substrate by plotting  $v_o/v_i$  vs. inhibitor concentration ( $[I]_o$ ) according to the rearranged Michaelis-Menten equation for competitive inhibition kinetics:  $v_o/v_i = 1 + [I]_o / (K_i (1 + [S]_o / K_m))$ , where  $v_o$  is the uninhibited initial velocity,  $v_i$  is the initial velocity in the presence of inhibitor at any given inhibitor concentration ( $[I]_o$ ) and  $[S]_o$  is the substrate concentration used. The resulting data were fitted using linear regression and the resulting slope,  $1/(K_i(1+[S]_o/K_m))$ , was used to calculate the  $K_i^*$  value.

The obtained  $K_i^*$  values for the various compounds of the present invention are given in the afore-mentioned Table wherein the compounds have been arranged in the order of ranges of  $K_i^*$  values. From these test results, it would be apparent to the skilled artisan that the compounds of the invention have excellent utility as NS3-serine protease inhibitors.

While the present invention has been described with in conjunction with the specific embodiments set forth above, many alternatives, modifications and other variations thereof will be apparent to those of ordinary skill in the art. All such alternatives, modifications and variations are intended to fall within the spirit and scope of the present invention.



Table 2







Compound From Example #	STRUCTURE	FORMULA	LRMS (FAB) M+H
1		C44 H68 N6 O8	809.5
2		C40 H60 N6 O8	753.5
3		C42 H62 N6 O8	779.5
4		C39 H54 N6 O8	735.4
5		C35 H54 N6 O8	687.4
6		C38 H51 Br N6 O8	720.4

Table 2

137








7		C38 H47 F5 N6 O8	811.3
8		C41 H53 N7 O10	804.4
9		C42 H54 N6 O8	771.4
10		C38 H49 F3 N6 O8	775.4
11		C35 H53 N5 O9	688.4
12		C39 H51 F3 N6 O8	789.4
13		C42 H57 N5 O9	776.4

Table 2

138








14		C34 H49 F3 N6 O8	727.4
15		C47 H59 N5 O10	854.4
16		C40 H53 N5 O10	764.4
17		C48 H61 N5 O10	868.4
18		C41 H55 N5 O10	778.4
19		C33 H49 N5 O9 S	692.3
20		C42 H59 N5 O9	778.4

Table 2

139








21		C38 H50 F N5 O9	740.4
22		C38 H50 Cl N5 O9	756.3
23		C38 H50 F N5 O9	740.4
24		C45 H61 N5 O10	832.4
25		C38 H57 N5 O10	744.4
26		C47 H68 N6 O9	861.5
27		C43 H68 N6 O9	813.5

Table 2

140

28		C41 H56 N6 O10	793.4
29		C37 H57 N5 O9	716.4
30		C39 H55 N5 O10	754.4
31		C46 H59 N5 O10	842.4
32		C36 H55 N5 O9	702.4
33		C40 H63 N5 O9	758.5
34		C34 H49 N5 O9	672.4

Table 2

141















35		C38 H57 N5 O9	728.4
36		C56 H71 N5 O9 S	990.5
37		C44 H55 N5 O9	798.4
38		C35 H49 N7 O9	712.4
39		C40 H61 N5 O10 S2	836.4
40		C39 H59 N5 O9 S2	806.4
41		C36 H53 N5 O9 S2	764.3

Table 2

142

42		C40 H61 N5 O9 S2	820.4
43		C35 H51 N5 O9 S2	750.3
44		C40 H64 N6 O11 Si	805.5
45		C34 H52 N6 O9	689.4
47		C37 H46 Cl2 N6 O10	805.3
48		C36 H54 F N5 O9	720.4
49		C35 H52 F N5 O9	706.4

143

50		C49 H62 N6 O11	911.5
51		C41 H56 N6 O10	793.4
52		C42 H56 N6 O12	837.4
53		C41 H57 N7 O12 S	872.4
54		C36 H54 N6 O10	731.4
55		C40 H62 N6 O10	787.5
56		C42 H64 N6 O10	813.5



Table 2

144








57		C40 H60 N6 O10	785.4
58		C38 H57 N7 O11	788.4
59		C41 H65 N7 O12 S	880.4
60		C40 H61 N5 O11	788.4
61		C39 H59 N5 O9	742.4
62		C35 H51 N5 O9	686.4
63		C41 H55 N5 O9 S2	826.4

Table 2

145








64		C36 H53 N5 O11	732.4
65		C39 H59 N5 O11	774.4
66		C35 H51 N5 O11	718.4
67		C45 H53 N5 O9	808.4
68		C38 H47 N5 O9	718.3
69		C39 H61 N5 O9	744.5
70		C37 H50 N6 O8	707.4

Table 2

146








71		C37 H49 N5 O9	708.4
72		C34 H51 N5 O9	674.4
73		C36 H56 N6 O10 S	765.4
74		C36 H48 N6 O10 S	757.3
75		C37 H47 N5 O10	722.3
76		C37 H45 Cl2 N5 O10	790.3
77		C34 H51 N5 O10 S	722.3

Table 2

147








78		C36 H56 N6 O11 S2	813.4
79		C36 H48 N6 O11 S2	805.3
80		C37 H47 N5 O11 S	770.3
81		C37 H45 Cl2 N5 O11 S	838.2
82		C39 H49 Cl2 N5 O10	818.3
83		C37 H57 N5 O8	700.4
84		C38 H60 N6 O10 S	793.4

Table 2

148








85		C38 H52 N6 O10 S	785.4
86		C39 H47 F4 N5 O10	822.3
87		C33 H51 N5 O9 S	694.3
88		C32 H49 N5 O9 S	680.3
89		C40 H63 N5 O11 S	822.4
90		C39 H55 N5 O9 S	770.4
91		C36 H55 N5 O11 S	766.4

Table 2

149








92		C38 H57 N5 O10	744.4
93		C34 H51 N5 O11	706.4
94		C40 H59 N5 O9	754.4
95		C38 H55 N5 O9	726.4
96		C36 H51 N5 O9	698.4
97		C34 H47 N5 O9	670.3
98		C42 H57 N5 O10	792.4

Table 2

150






















99		2 C40 H61 N5 O10	772.4
100		C37 H48 F N5 O8	710.4
101		C36 H49 N5 O8 S	712.3
102		C40 H54 Cl N5 O8	768.4
103		C39 H57 N5 O9	740.4
104		C35 H49 N5 O9	684.4
105		C37 H53 N5 O9	712.4

Table 2

151

106		C36 H52 N6 O8	697.4
107		C48 H66 N8 O10	915.5
108		C44 H58 N8 O10	859.4
109		C48 H65 N5 O10	872.5
110		C39 H49 N5 O9	732.4
111		C37 H53 N5 O9	712.4
112		C33 H45 N5 O9	656.3



113		C38 H56 N6 O9	741.4
114		C37 H54 N6 O9	727.4
115		C41 H61 N5 O9	768.5
116		C37 H53 N5 O9	712.4
117		C39 H58 N6 O8	739.4
118		C40 H60 N6 O8	753.5
119		C39 H58 N6 O9	755.4








120		C38 H56 N6 O8	725.4
121		C38 H57 N5 O9	728.4
122		C34 H49 N5 O9	672.4
123		C40 H55 N5 O9	750.4
124		C39 H53 N5 O9	736.4
125		C39 H53 N5 O9	736.4
126		C35 H45 N5 O9	680.3

Table 2

154






















127		C36 H47 N5 O9	694.3
128		C38 H60 N6 O8	729.5
129		C41 H58 N6 O8	763.4
130		C36 H54 N6 O8	699.4
131		C43 H57 N5 O8	772.4
132		C39 H49 N5 O8	716.4
133		C42 H55 N5 O9	774.4

Table 3

STRUCTURE	KI* CLASS
	B
	B
	B
	B
	B
	B
	B
	B

STRUCTURE	KI* CLASS
	B
	B
	B
	C
	C
	B








STRUCTURE	NAME	Ki* Range
	(2-CO2,3-Me)PhCO-G(Chx)-L-nV-(CO)-G-G(Ph)-Am	a
	(2,5-diF,6-CO2)PhCO-G(Chx)-L-nV-(CO)-G-G(Ph)-Am	a
	(2-CO2)BnCO-G(Chx)-L-nV-(CO)-G-G(Ph)-Am	c
	(2-SO3)PhCO-G(Chx)-L-nV-(CO)-G-G(Ph)-Am	b
	(2-CO2)cyclopentenoyl-G(Chx)-L-nV-(CO)-G-G(Ph)-Am	c
	(2-CO2,3-OH)PhCO-G(Chx)-L-nV-(CO)-G-G(Ph)-Am	a
	(2,3,4,5-tetraF,6-CO2)PhCO-G(Chx)-L-nV-(CO)-G-G(Ph)-Am	a



Table 4

158

STRUCTURE	NAME	Ki* Range
	(2CO <sub>2</sub> )PhCO- G(Chx)-C((O <sub>2</sub> )Et)- nV-(CO)-G-G(Ph)- Am	b
	(2CO <sub>2</sub> )PhCO- G(Chx)-C((O <sub>2</sub> )Ph)- nV-(CO)-G-G(Ph)- Am	b
	(2CO <sub>2</sub> )PhCO- G(Chx)- C((O <sub>2</sub> )EtPh)-nV- (CO)-G-G(Ph)-Am	b
	iBoc-G(Chx)- C((O <sub>2</sub> )Et)-nV-(CO)- G-G(Ph)-Am	b
	(2CO <sub>2</sub> )PhCO- G(Chx)-C((O <sub>2</sub> )- 1Np)-nV-(CO)-G- G(Ph)-Am	b
	(2CO <sub>2</sub> )PhCO- G(Chx)-C((O <sub>2</sub> )- 2Np)-nV-(CO)-G- G(Ph)-Am	b
	iBoc-G(Chx)- C((O <sub>2</sub> )Ph)-nV-(CO)- G-G(Ph)-Am	b

Table 4

159

STRUCTURE	NAME	Ki* Range
	((3-tetrahydrofuran)-CH2O)CO-G(Chx)-L-nV-(CO)-G-G(Ph)-Am	b
	7-Me-2-iPr-Oct-4-ene-(CO)-L-nV-(CO)-G-G(Ph)-OH	c




160

Claims

What is claimed is:

1. A compound, including enantiomers, stereoisomers, rotamers and tautomers of said compound, and pharmaceutically acceptable salts or solvates of said compound, said compound having the general structure shown in Formula I:

Formula I

wherein:

- G, J and Y may be the same or different and are independently selected from the group consisting of the moieties: H, alkyl, alkyl-aryl, heteroalkyl, heteroaryl, aryl-heteroaryl, alkyl-heteroaryl, cycloalkyl, alkyloxy, alkyl-aryloxy, aryloxy, heteroaryloxy, heterocycloalkyloxy, cycloalkyloxy, alkylamino, arylamino, alkyl-aryl amino, arylamino, heteroaryl amino, cycloalkyl amino and heterocycloalkyl amino, with the proviso that Y may be additionally optionally substituted with  $X^{11}$  or  $X^{12}$ ;

- $X^{11}$  is selected from the group consisting of alkyl, alkenyl, alkynyl, cycloalkyl, cycloalkyl-alkyl, heterocyclyl, heterocyclylalkyl, aryl, alkylaryl, arylalkyl, heteroaryl, alkylheteroaryl, or heteroarylalkyl moiety, with the proviso that  $X^{11}$  may be additionally optionally substituted with  $X^{12}$ ;

- $X^{12}$  is hydroxy, alkoxy, aryloxy, thio, alkylthio, arylthio, amino, alkylamino, arylamino, alkylsulfonyl, arylsulfonyl, alkylsulfonamido, arylsulfonamido, carboxy, carbalkoxy, carboxamido, alkoxy carbonyl amino, alkoxy carbonyloxy, alkylureido, arylureido, halogen, cyano, or nitro, with the

proviso that said alkyl, alkoxy, and aryl may be additionally optionally substituted with moieties independently selected from  $X^{12}$ ;

$R^1$  is  $COR^5$  or  $B(OR)_2$ , wherein  $R^5$  is selected from the group consisting of H, OH,  $OR^8$ ,  $NR^9R^{10}$ ,  $CF_3$ ,  $C_2F_5$ ,  $C_3F_7$ ,  $CF_2R^6$ ,  $R^6$  and

- 5  $COR^7$  wherein  $R^7$  is selected from the group consisting of H, OH,  $OR^8$ ,  $CHR^9R^{10}$ , and  $NR^9R^{10}$ , wherein  $R^6$ ,  $R^8$ ,  $R^9$  and  $R^{10}$  may be the same or different and are independently selected from the group consisting of H, alkyl, aryl, heteroalkyl, heteroaryl, cycloalkyl, cycloalkyl, arylalkyl, heteroarylalkyl,  $CH(R^{1'})COOR^{11}$ ,  $CH(R^{1'})CONR^{12}R^{13}$ ,  $CH(R^{1'})CONHCH(R^{2'})COOR^{11}$ ,  
 10  $CH(R^{1'})CONHCH(R^{2'})CONR^{12}R^{13}$ ,  $CH(R^{1'})CONHCH(R^{2'})R'$ ,  $CH(R^{1'})CONHCH(R^{2'})CONHCH(R^{3'})COOR^{11}$ ,  $CH(R^{1'})CONHCH(R^{2'})CONHCH(R^{3'})CONR^{12}R^{13}$ ,  $CH(R^{1'})CONHCH(R^{2'})CONHCH(R^{3'})CONHCH(R^{4'})COOR^{11}$ ,  $CH(R^{1'})CONHCH(R^{2'})CONHCH(R^{3'})CONHCH(R^{4'})CONR^{12}R^{13}$ ,  
 15  $CH(R^{1'})CONHCH(R^{2'})CONHCH(R^{3'})CONHCH(R^{4'})CONHCH(R^{5'})COOR^{11}$ , and  $CH(R^{1'})CONHCH(R^{2'})CONHCH(R^{3'})CONHCH(R^{4'})CONHCH(R^{5'})CONR^{12}R^{13}$ , wherein  $R^{1'}$ ,  $R^{2'}$ ,  $R^{3'}$ ,  $R^{4'}$ ,  $R^{5'}$ ,  $R^{11}$ ,  $R^{12}$ ,  $R^{13}$ , and  $R'$  may be the same or different and are independently selected from a group consisting of H, alkyl,  
 20 aryl, heteroalkyl, heteroaryl, cycloalkyl, alkyl-aryl, alkyl-heteroaryl, aryl-alkyl and heteroaralkyl;

Z is selected from O, N, or CH;

W maybe present or absent, and if W is present, W is selected from  $C=O$ ,  $C=S$ , or  $SO_2$ ; and

- 25 R,  $R^2$ ,  $R^3$  and  $R^4$  are independently selected from the group consisting of H; C1-C10 alkyl; C2-C10 alkenyl; C3-C8 cycloalkyl; C3-C8 heterocycloalkyl, alkoxy, aryloxy, alkylthio, arylthio, amino, amido, ester,

carboxylic acid, carbamate, urea, ketone, aldehyde, cyano, nitro; oxygen, nitrogen, sulfur, or phosphorus atoms (with said oxygen, nitrogen, sulfur, or phosphorus atoms numbering zero to six); (cycloalkyl)alkyl and (heterocycloalkyl)alkyl, wherein said cycloalkyl is made of three to eight  
 5 carbon atoms, and zero to six oxygen, nitrogen, sulfur, or phosphorus atoms, and said alkyl is of one to six carbon atoms; aryl; heteroaryl; alkyl-aryl; and alkyl-heteroaryl;

wherein said alkyl, heteroalkyl, alkenyl, heteroalkenyl, aryl, heteroaryl, cycloalkyl and heterocycloalkyl moieties may be optionally substituted, with  
 10 said term "substituted" referring to optional and chemically-suitable substitution with one or more moieties selected from the group consisting of alkyl, alkenyl, alkynyl, aryl, aralkyl, cycloalkyl, heterocyclic, halogen, hydroxy, thio, alkoxy, aryloxy, alkylthio, arylthio, amino, amido, ester, carboxylic acid, carbamate, urea, ketone, aldehyde, cyano, nitro, sulfonamide, sulfoxide,  
 15 sulfone, sulfonylurea, hydrazide, and hydroxamate.

2. The compound of claim 1, wherein  $R^1$  is  $COR^5$ , and  $R^5$  is H, OH,  $COOR^8$  or  $CONR^9R^{10}$ .

3. The compound of claim 2, wherein  $R^1$  is  $COCONR^9R^{10}$ , and is  $R^9$  is H,  $R^{10}$  is selected from the group consisting of H,  $CH(R^{1'})COOR^{11}$ ,  $CH(R^{1'})$   
 20  $CONR^{12}R^{13}$ ,  $CH(R^{1'})CONHCH(R^{2'})COOR^{11}$ ,  $CH(R^{1'})CONHCH(R^{2'})$   
 $CONR^{12}R^{13}$ , and  $CH(R^{1'})CONHCH(R^{2'})(R')$ .

4. The compound of claim 3, wherein  $R^{10}$  is  
 $CH(R^{1'})CONHCH(R^{2'})COOR^{11}$ ,  $CH(R^{1'})CONHCH(R^{2'})CONR^{12}R^{13}$ , or  
 $CH(R^{1'})CONHCH(R^{2'})(R')$ , wherein  $R^{1'}$  is H or alkyl, heteroalkyl and  $R^{2'}$  is  
 25 phenyl, substituted phenyl, hetero atom-substituted phenyl, thiophenyl, cycloalkyl, piperidyl and pyridyl.


5. The compound of claim 4, wherein  $R^{1'}$  is H.

6. The compound of claim 5, wherein

$R^{11}$  is H or *tert*-butyl;

R' is hydroxymethyl; and

R<sup>2'</sup> is selected from the group consisting of:



5

wherein:

U<sup>1</sup> and U<sup>2</sup> maybe same or different and are independently selected from the group consisting of H, F, CH<sub>2</sub>COOH, CH<sub>2</sub>COOMe, CH<sub>2</sub>CONH<sub>2</sub>, CH<sub>2</sub>CONHMe, CH<sub>2</sub>CONMe<sub>2</sub>, azido, amino, hydroxyl, substituted amino, substituted hydroxyl;

10


U<sup>3</sup> and U<sup>4</sup> maybe same or different and are O or S;

U<sup>5</sup> is selected from the moieties consisting of alkylsulfonyl, aryl sulfonyl, heteroalkyl sulfonyl, heteroaryl sulfonyl, alkyl carbonyl, aryl carbonyl, heteroalkyl carbonyl, heteroaryl carbonyl, alkoxy carbonyl, aryloxy carbonyl, heteroaryloxy carbonyl, alkylaminocarbonyl, arylaminocarbonyl and heteroarylamino carbonyl or combinations thereof; and

15


NR<sup>12</sup>R<sup>13</sup> is selected from the group consisting of:

164

























 $\text{NH}_2, \text{NHMe}, \text{N(Me)OMe}, \text{NMe}_2,$ 

wherein  $U^6$  is H, OH, or  $CH_2OH$ .

7. The compound of claim 2, wherein R<sup>2</sup> is selected from the group consisting of the following moieties:




8. The compound of claim 7, wherein  $R^3$  is selected from the group consisting of:




166

and




wherein  $R^{31} = \text{OH}$  or O-alkyl;

$Y^{19}$  is selected from the following moieties:




5

and  $Y^{20}$  is selected from the following moieties:



9. The compound of claim 8, wherein  $R^3$  is selected from the following structures:


167







































10. The compound of claim 9, wherein  $Z = N$  and  $R^4 = H$ .
11. A compound of claim 10, wherein  $W$  is  $C=O$ , or  $SO_2$ .
12. A compound of claim 11, wherein  $Y$  is selected from the following  
5 moieties:




168




169



170



171




wherein:

Y<sup>11</sup> is selected from H, COOH, COOEt, Ome, Ph, Oph, NHMe, NHAc, NHPh, CH(Me)<sub>2</sub>, 1-triazolyl, 1-imidazolyl, and NHCH<sub>2</sub>COOH;

Y<sup>12</sup> is selected from H, COOH, COOMe, Ome, F, Cl, or Br;

$Y^{13}$  is selected from the following moieties:



Y<sup>14</sup> is selected from MeSO<sub>2</sub>, Ac, Boc, *t*Boc, Cbz, or Alloc;

172


$Y^{15}$  and  $Y^{16}$  may be the same or different and are independently selected from alkyl, aryl or heteroalkyl, or heteroaryl;

$Y^{17}$  is  $CF_3$ ,  $NO_2$ ,  $CONH_2$ ,  $OH$ ,  $COOCH_3$ ,  $OCH_3$ ,  $OC_6H_5$ ,  $C_6H_5$ ,  $COC_6H_5$ ,  $NH_2$ , or  $COOH$ ; and

5  $Y^{18}$  is  $COOCH_3$ ,  $NO_2$ ,  $N(CH_3)_2$ ,  $F$ ,  $OCH_3$ ,  $CH_2COOH$ ,  $COOH$ ,  $SO_2NH_2$ , or  $NHCOCH_3$ .

13. A compound of claim 12, wherein Y is selected from the group consisting of:

173




wherein:

Y<sup>17</sup> = CF<sub>3</sub>, NO<sub>2</sub>, CONH<sub>2</sub>, OH, NH<sub>2</sub>, or COOH;Y<sup>18</sup> = F, COOH,


174

14. The compound of claim 13, wherein J is selected from the group consisting of:




15. The compound of claim 14 where in J is H, CH<sub>3</sub> or Bn.
16. The compound of claim 15 wherein G is selected from moieties:













175








Q=







- 5 17. The compound of claim 16, wherein G is selected from the group consisting of :





18. A pharmaceutical composition comprising as an active ingredient a compound of claim 1.
19. The pharmaceutical composition of claim 18 for use in treating disorders associated with Hepatitis C virus.
20. The pharmaceutical composition of claim 18 additionally comprising a pharmaceutically acceptable carrier.
21. A method of treating disorders associated with the HCV protease, said method comprising administering to a patient in need of such treatment a pharmaceutical composition which composition comprises therapeutically effective amounts of a compound of claim 1.
22. The method of claim 21, wherein said administration is via subcutaneous administration.
23. The use of a compound of claim 1 for the manufacture of a medicament to treat disorders associated with the HCV protease.


24. A method of preparing a pharmaceutical composition for treating disorders associated with the HCV protease, said method comprising bringing into intimate contact a compound of claim 1 and a pharmaceutically acceptable carrier.
25. A compound exhibiting HCV protease inhibitory activity, including enantiomers, stereoisomers, rotamers and tautomers of said compound, and pharmaceutically acceptable salts or solvates of said compound, said compound being selected from the group of compounds with structures listed below:



179




5



R = Me

R = Benzyl



10

X = O<sup>+</sup>Bu


X = OH

X = NH<sub>2</sub>


X = NMeOMe

X = NMe<sub>2</sub>


180

X = O<sup>t</sup>Bu


X = OH



5




X = H, Y = tBu; X = tBu, Y = H





10

R = Propargyl; R = Allyl


X = O<sup>t</sup>Bu; X = OH

181


X = O<sup>t</sup>Butyl

X = OH

5

X = O<sup>t</sup>Butyl


X = OH

X = NMe<sub>2</sub>X = O<sup>t</sup>Bu

X = OH


10

182

R = <sup>t</sup>Bu

R = H




R = Me




X = H, Y = COOH


X = COOH, Y = H

5








26. A pharmaceutical composition for treating disorders associated with the HCV protease, said composition comprising therapeutically effective amount of one or more compounds in claim 25 and a pharmaceutically acceptable carrier.
27. The pharmaceutical composition of claim 26, additionally containing an antiviral agent.
28. The pharmaceutical composition of claim 26 or claim 27, still additionally containing an interferon.
29. The pharmaceutical composition of claim 28, wherein said antiviral agent is ribavirin and said interferon is  $\alpha$ -interferon.
30. A compound selected from the group consisting of:











5





10



15




20



25

and



30

or an enantiomer, stereoisomer, rotamer or tautomer thereof, or a pharmaceutically acceptable salt or solvate thereof, wherein the compound exhibits HCV inhibitory activity.

- 5 31. A pharmaceutical composition, comprising one or more compounds of claim 30.
32. A method of treatment of an hepatitis C virus associated disorder, comprising administering an effective amount of one or more compounds of claim 30.
- 10 33. A method of modulating the activity of hepatitis C virus (HCV) protease, comprising contacting HCV protease with one or more compounds of claim 30.
34. A method of treating, preventing, or ameliorating one or more symptoms of hepatitis C, comprising administering an effective amount of one  
15 or more compounds of claim 30.
35. The method of claim 33, wherein the HCV protease is the NS3/NS4a protease.
36. The method of claim 35, wherein the compound or compounds inhibit HCV NS3/NS4a protease.
- 20 37. A method of modulating the processing of hepatitis C virus (HCV) polypeptide, comprising contacting a composition containing the HCV polypeptide under conditions in which the polypeptide is processed with one or more compounds of claim 30.
38. The compound of claim 7, wherein R<sup>3</sup> is cyclohexyl.
- 25 39. The compound of claim 11, wherein Y is selected from the group consisting of 2-carboxy-3-hydroxyphenyl, 3-tetrahydrofurylmethoxy, and 2-sulfophenyl.
40. The compound of claim 15, wherein G is selected from the group consisting of ethylsulfonylmethyl, phenylsulfonylmethyl, 2-  
30 phenylethylsulfonylmethyl and 1-naphthylsulfonylmethyl.

## INTERNATIONAL SEARCH REPORT

International application No.

PCT/US01/22813

## A. CLASSIFICATION OF SUBJECT MATTER

IPC(7) : Please See Extra Sheet.

US CL : Please See Extra Sheet.

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 546/233, 234; 548/336.1, 452; 549/21, 22, 35, 39, 357, 430; 560/27, 37, 38, 39, 40, 41; 562/442, 443, 444, 445; 564/153

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

NONE

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

CASONLINE

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 5,514,694 A (POWERS et al) 07 May 1996	1-40
A	US 5,488,067 A (HANSON) 30 January 1996	1-40
A	WO 98/22496 A2 (F. HOFFMANN-LA ROCHE AG) 28 May 1998.	1-40



Further documents are listed in the continuation of Box C.



See patent family annex.

\*

Special categories of cited documents:

\*T\*

later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*A\*

document defining the general state of the art which is not considered to be of particular relevance

\*X\*

document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*B\*

earlier document published on or after the international filing date

\*Y\*

document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

\*L\*

document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\*

document referring to an oral disclosure, use, exhibition or other means

\*P\*

document published prior to the international filing date but later than the priority date claimed

\*Q\*

document member of the same patent family

Date of the actual completion of the international search

25 SEPTEMBER 2001

Date of mailing of the international search report

29 OCT 2001

Name and mailing address of the ISA/US

Commissioner of Patents and Trademarks

Box PCT

Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

SHAILENDRA KUMAR

Telephone No. (703) 308-1235

INTERNATIONAL SEARCH REPORT

International application No.  
PCT/US01/22813

A. CLASSIFICATION OF SUBJECT MATTER:  
IPC (7):

C07D 209/02, 211/04, 233/56, 317/10, 319/04, 339/02, 339/08; C07C 229/00, 233/05, 271/08, 271/32; A61K 31/16, 31/27/31/195, 31/357, 31/385, 31/403, 31/445, 31/4164

A. CLASSIFICATION OF SUBJECT MATTER:  
US CL :

546/233, 234; 548/336.1, 452; 549/21, 22, 35, 39, 357, 430; 560/27, 37, 38, 39, 40, 41; 562/442, 443, 444, 445; 564/153